

## **SECRETED HUMAN PROTEINS**

5 This application claims the benefit of copending provisional application  
Serial No. 60/032,757, filed December 11, 1996, which is incorporated herein by  
reference.

### **TECHNICAL AREA OF THE INVENTION**

10 The invention relates to the area of proteins. More particularly, the  
invention relates to human secreted proteins.

### **BACKGROUND OF THE INVENTION**

15 Secreted proteins include such important proteins as growth factors,  
cytokines and their receptors, extracellular matrix proteins, and proteases.

Nucleotide sequences encoding these proteins can be used to detect disease states in  
which such proteins are implicated and to develop therapeutics for such diseases.

Thus, there is a need in the art for methods of identifying secreted proteins and the  
nucleotide sequences which encode them.

### **SUMMARY OF THE INVENTION**

20 It is an object of the invention to provide an isolated and purified human  
protein.

It is yet another object of the invention to provide a fusion protein.

It is still another object of the invention to provide a preparation of antibodies.

It is even another object of the invention to provide an isolated and purified subgenomic polynucleotide.

5 It is yet another object of the invention to provide an isolated gene.

It is a further object of the invention to provide a DNA construct for expressing all or a portion of a human protein.

It is still another object of the invention to provide a host cell comprising a DNA construct.

10 It is another object of the invention to provide a homologously recombinant cell.

It is even another object of the invention to provide a method of producing a human protein.

15 It is another object of the invention to provide a method of identifying a secreted polypeptide which is modified by rough microsomes.

These and other objects of the invention are provided by one or more of the embodiments described below.

20 One embodiment of the invention provides an isolated and purified human protein. The isolated and purified human protein has an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

25 Another embodiment of the invention provides an isolated and purified human protein having an amino acid sequence which is at least 85% identical to an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

30 Still another embodiment of the invention provides a polypeptide comprising at least 6 contiguous amino acids of an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

Even another embodiment of the invention provides a fusion protein. The fusion protein comprises a first protein segment and a second protein segment fused together by means of a peptide bond. The first protein segment consists of at least 6 contiguous amino acids selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

Yet another embodiment of the invention provides a preparation of antibodies. The antibodies specifically bind to a human protein having an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

Even another embodiment of the invention provides an isolated and purified subgenomic polynucleotide. The isolated and purified subgenomic polynucleotide has a nucleotide sequence selected from the group consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

Yet another embodiment of the invention provides an isolated and purified subgenomic polynucleotide consisting of at least 10 contiguous nucleotides selected from the group consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

Still another embodiment of the invention provides an isolated gene. The isolated gene corresponds to a cDNA sequence selected from the group consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

Another embodiment of the invention provides a DNA construct for expressing all or a portion of a human protein. The DNA construct comprises a promoter and a polynucleotide segment. The polynucleotide segment encodes at least 6 contiguous amino acids of a human protein having an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

The polynucleotide segment is located downstream from the promoter.

Transcription of the polynucleotide segment initiates at the promoter.

Even another embodiment of the invention provides a host cell comprising a DNA construct. The DNA construct comprises a promoter and a polynucleotide segment. The polynucleotide segment encodes at least 6 contiguous amino acids of a human protein having an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38. The polynucleotide segment is located downstream from the promoter. Transcription of the polynucleotide segment initiates at the promoter.

Still another embodiment of the invention provides a homologously recombinant cell having incorporated therein a new transcription initiation unit. The transcription initiation unit comprises in 5' to 3' order an exogenous regulatory sequence, an exogenous exon, and a splice donor site. The transcription initiation unit is located upstream to a coding sequence of a gene. The gene comprises a nucleotide sequence selected from the group consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19. The exogenous regulatory sequence controls transcription of the coding sequence of the gene.

Yet another embodiment of the invention provides a method of producing a human protein. A culture of a cell is grown. The cell comprises a DNA construct. The DNA construct comprises a promoter and a polynucleotide segment. The polynucleotide segment encodes at least 6 contiguous amino acids of a human protein having an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38. The polynucleotide segment is located downstream from the promoter. Transcription of the polynucleotide segment initiates at the promoter. The protein is purified from the culture.

Even another embodiment of the invention provides a method of producing a human protein. A culture of a cell is grown. The cell comprises a new transcription initiation unit. The transcription initiation unit comprises in 5' to 3'

order an exogenous regulatory sequence, an exogenous exon, and a splice donor site. The transcription initiation unit is located upstream to a coding sequence of a gene. The gene comprises a nucleotide sequence selected from the group consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19. The exogenous regulatory sequence controls transcription of the coding sequence of the gene. The protein is purified from the culture.

Another embodiment of the invention provides a method of identifying a secreted polypeptide which is modified by rough microsomes. A population of cDNA molecules is transcribed *in vitro* whereby a population of cRNA molecules is formed. A first portion of the population of cRNA molecules is translated *in vitro* in the absence of rough microsomes whereby a first population of polypeptides is formed. A second portion of the population of cRNA molecules is translated *in vitro* in the presence of rough microsomes whereby a second population of polypeptides is formed. The first population of polypeptides is compared with the second population of polypeptides. Polypeptide members of the second population which have been modified by the rough microsomes are detected.

The present invention thus provides the art with a method for identifying secreted proteins or polypeptides, the amino acid sequences of nineteen novel human secreted proteins, and the nucleotide sequences which encode these proteins. The invention can be used to, *inter alia*, to produce secreted proteins for therapeutic and diagnostic purposes.

#### **DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS**

The inventors have discovered a method for identifying secreted proteins or polypeptides. Secreted proteins or polypeptides include soluble proteins which can be transported across a membrane, such as a cell membrane, nuclear membrane, or membrane of the endoplasmic reticulum, as well as proteins which can be partially secreted from a cell, such as membrane-bound receptors.

Secreted proteins can contain a signal (or secretion leader) sequence, located at the N-terminus and including at least several hydrophobic amino acids,

such as phenylalanine, methionine, leucine, valine, or tryptophan. Non-hydrophobic amino acids can also be included in the signal sequence. Signal sequences are described in von Heijne, *J. Mol. Biol.* 184:99-105 (1985) and Kaiser and Botstein, *Mol. Cell. Biol.* 6:2382-2391 (1986). Secreted proteins can also be glycosylated by post-translational modification. The presence of a signal sequence or the presence of glycosylation or both indicate that a particular protein is a secreted protein.

In order to identify secreted proteins or polypeptides, the method of the invention exploits properties of microsomes, which are the closed vesicles that result from fragmentation of endoplasmic reticulum. Microsomes can be rough or smooth, depending on whether the endoplasmic reticulum from which they were derived is studded with ribosomes. Microsomes, particularly rough microsomes, have the ability to perform post-translational modifications, such as glycosylation and cleavage of signal sequences from proteins or polypeptides.

To identify secreted proteins, a population of complementary DNA (cDNA) molecules is transcribed *in vitro* to synthesize a population of complementary RNA (cRNA) molecules. The cDNA molecules can be synthesized by reverse transcription of mRNA molecules isolated from a particular cell or tissue type or organism using, for example, a commercially available reverse transcriptase enzyme. Alternatively, the reverse transcription reaction to form cDNA molecules can be conducted on total RNA, without a preliminary purification of mRNA.

Any organism, such as a bacterium, plant, invertebrate, or vertebrate organism, can be used as a source of RNA. Particularly preferred sources of RNA are mammals, most preferably humans. Tissues, such as liver, brain, kidney, spleen, pancreas, or muscle, can be used as a source of RNA. Individual cell types, either primary cells or members of established cell lines, such as HeLa, CHO, PC12, P19, BHK, COS, or HepG2, are suitable sources of RNA. Tissues or primary cells isolated from organisms at a particular stage in development can be used as RNA sources. Stem cells, such as hematopoietic, neuronal, and embryonic stem cells, can also be used as a source of RNA.

Total RNA or mRNA can be isolated using methods known in the art. Such methods are described, *inter alia*, in Sambrook *et al.*, MOLECULAR CLONING, A

LABORATORY MANUAL (2d ed., Cold Spring Harbor Press, N.Y., 1989), and Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY (Greene Publishing Associates and John Wiley & Sons, N.Y., 1994). Techniques for RNA isolation can be tailored for a particular organism or cell type, as is known in the art.

5 Complementary DNA can optionally be obtained from a cDNA library. The cDNA library can be derived from the genome of any organism of interest, particularly a mammal or a human. Tissue- or cell type-specific cDNA libraries can also be used as a source of cDNA.

10 Transcription of cDNA molecules *in vitro* to form cRNA molecules can be carried out using any methods known in the art. These methods include, for example, placing cDNA into a cloning vector containing a promoter, such as an SP6, T7, or T3 polymerase promoter, and transcribing the cDNA using the appropriate polymerase. A variety of commercial kits are available for this purpose.

15 A first portion of the population of cRNA molecules can be translated *in vitro*, in the absence of rough microsomes, to form a first population of polypeptides which have not been post-translationally modified. A second portion of the population of cRNA molecules can be translated *in vitro* in the presence of rough microsomes. Under the conditions of the *in vitro* translation reaction, rough microsomes can cleave signal sequences from those polypeptides which comprise such sequences. Under the same conditions, rough microsomes can also glycosylate those polypeptides which contain glycosylation sites.

20 Methods of *in vitro* translation are those which are known in the art, such as translation in a reticulocyte lysate system, particularly a rabbit reticulocyte lysate. Reticulocyte lysate systems can be assembled in the laboratory or purchased  
25 \* commercially in kit form.

30 Microsomes can be prepared by disruption of tissues or cells by homogenization, as is known in the art. If desired, rough and smooth microsomes can be separated using well-known techniques, such as sucrose density gradient sedimentation. Microsomes are also available commercially, for example, such as the canine pancreatic microsomes available from Promega Corp., Madison, WI.

The first population of polypeptides can then be compared with the second population of polypeptides. This comparison can be by means of, for example, one- or two-dimensional polyacrylamide gel electrophoresis, as is known in the art. Polypeptides separated in the gels can be detected by any means known in the art, such as staining with copper, silver, Coomassie Brilliant Blue, amido black, fast green FCF, Ponceau S, or a chromophoric label. Separated proteins can also be visualized using radioactive, chemiluminescent, fluorescent, or enzymatic tags incorporated into the proteins before separation.

The gels can be dried or the proteins can be transferred to membranes, such as polyvinylidene difluoride membranes. Either the gels or membranes themselves or photographs of the gels or membranes can be compared by eye. Alternatively, the gels or membranes can be scanned, for example, with a densitometer and analyzed with the aid of a computer.

Polypeptide members of the second population of polypeptides, which have been modified by the rough microsomes, can be detected by any means available in the art. For example, a shift in the position of a polypeptide band can be observed, indicating an increase in molecular weight of a member of the second population compared with the corresponding polypeptide member of the first population. Such an increase in molecular weight indicates that the polypeptide member of the second population was glycosylated by the rough microsomes.

A shift in the position of a polypeptide band indicating a decrease in molecular weight of a member of the second population compared with the corresponding polypeptide member of the first population can also be observed. This decrease in molecular weight indicates that the polypeptide member of the second population contained a signal sequence which was cleaved by the rough microsomes.

Polypeptides which are modified by the rough microsomes are identified as secreted polypeptides. Optionally, quantities of cDNA molecules which encode secreted polypeptides can be obtained. Molecules of cDNA which encode polypeptides which are post-translationally modified by the rough microsomes can be placed into suitable vectors using standard recombinant DNA techniques and



used to transform host cells. Many vectors are available for this purpose, such as retroviral or adenoviral vectors and bacteriophage, as described below.

Vectors comprising cDNA which encode secreted polypeptides can be introduced into host cells using techniques available in the art. These techniques include, but are not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, and calcium phosphate-mediated transfection.

The host cells can be any host cells which are capable of propagating cDNA molecules. A variety of host cells, for example immortalized cell lines such as HeLa, CHO, or HEK, are available for this purpose.

Transformed host cells can be diluted serially and cultured to form individual colonies. Methods of culturing host cells and the media suitable for each host cell type are well known in the art. Preferably, each colony originates from a single transformed host cell. Separate preparations of cDNA from each colony can be prepared, as described above, and transcribed *in vitro* to form cRNA. The cRNA can be transcribed to form secreted polypeptides, which can be purified as is known in the art. If the preparation of secreted polypeptides from a colony contains more than one species of polypeptide, the steps described above can be repeated until a colony is obtained which contains cDNA encoding only a single species of polypeptide.

Complementary DNA molecules which encode secreted proteins can be sequenced using standard nucleotide sequencing techniques. The sequence of each cDNA molecule can be compared with known sequences in a database to determine whether the clone encodes a known or a novel secreted protein.

The inventors have used the method of the invention to identify nineteen novel human secreted proteins. Amino acid sequences for these nineteen human secreted proteins are disclosed in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38. Nucleotide sequences which encode the proteins are disclosed in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19, respectively.

Clones containing the cDNAs of the secreted proteins were deposited on December 11, 1997, with the ATCC. Individual bacterial cells (*E. coli*) in this composite deposit contain one or more of the polynucleotides encoding the secreted proteins of the invention and can be retrieved using an oligonucleotide probe designed from the sequence for that particular polynucleotide, as provided herein. Each polynucleotide can be removed from the vector by performing an EcoRI/NotI digestion (5' site, EcoRI; 3' site, NotI). The deposit submitted to the ATCC has been designated SECP120997. The nucleotide sequences of these deposits and the amino acid sequences they encode are controlling in the event of a discrepancy between the amino acid and nucleotide sequences disclosed herein and those contained in the deposits.

A purified and isolated subgenomic polynucleotide of the present invention comprises at least 10, 12, 15, 18, 20, 25, 30, 35, 40, 45, or 50 contiguous nucleotides selected from the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19. The isolated and purified subgenomic polynucleotides can comprise an entire nucleotide sequence selected from the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

Subgenomic polynucleotides contain less than a whole chromosome and are preferably intron-free. Polynucleotides of the invention can be isolated and purified free from other nucleotide sequences by standard nucleic acid purification techniques, using restriction enzymes and probes to isolate fragments comprising the coding sequences.

Isolated genes corresponding to the cDNA sequences disclosed herein are also provided. Known methods can be used to isolate the corresponding genes using the provided cDNA sequences. These methods include preparation of probes or primers from the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19 for use in identifying or amplifying the genes from human genomic libraries or other sources of human genomic DNA.

The coding sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19 can be made using reverse transcriptase with

human mRNA as a template. Amplification by PCR can also be used to obtain the polynucleotides, using either genomic DNA or cDNA as a template. Polynucleotide molecules of the invention can also be made using the techniques of synthetic chemistry given the sequences disclosed herein. The degeneracy of the genetic code permits alternate nucleotide sequences which will encode the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38 to be synthesized. All such nucleotide sequences are within the scope of the present invention.

Polynucleotide molecules of the invention can be propagated in vectors and cell lines as is known in the art. Polynucleotide molecules can be on linear or circular molecules. They can be on autonomously replicating molecules or on molecules without replication sequences. For propagation, polynucleotides of the invention can be introduced into suitable host cells using any techniques available in the art, as described above.

Subgenomic polynucleotides of the invention can be used to propagate additional copies of the polynucleotides or to express protein, polypeptides, or fusion proteins. The subgenomic polynucleotides disclosed herein can also be used, for example, as biomarkers for tissues or chromosomes, as molecular weight markers for DNA gels, to elicit immune responses, such as the formation of antibodies against single- or double-stranded DNA, and in DNA-ligand interaction assays, to detect proteins or other molecules which interact with the nucleotide sequences.

Disease states may be associated with alterations in the expression of genes which encode proteins of the invention. Polynucleotide sequences disclosed herein can also be used to determine the involvement of any of these sequences in disease states. For example, a gene in a diseased cell can be sequenced and compared with a wild-type coding sequence of the invention. Alternatively, nucleotide probes can be constructed and used to detect normal or altered (mutant) forms of mRNA in a diseased cell. Subgenomic polynucleotides of the invention can also be used to design diagnostic tests and therapeutic compositions for diseases which may be associated with altered expression of these genes.

The present invention provides both full-length and mature forms of the disclosed proteins. Full-length forms of the proteins have the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38. The full-length forms of a protein can be processed enzymatically to remove a signal sequence, resulting in a mature form of the protein. Signal sequences can be identified by examination of the amino acid sequences disclosed herein and comparison with amino acid sequences of known signal sequences (see, *e.g.*, von Heijne, 1985; Kaiser & Botstein, 1986). Similarly, transmembrane domains can be identified by examination of the amino acid sequences disclosed herein. A transmembrane domain typically contains a long stretch of 15-30 hydrophobic amino acids.

Other domains with predicted functions can also be identified. For example, the protein having the amino acid sequence shown in SEQ ID NO:23 comprises a Kunitz type serine protease inhibitor domain spanning amino acids 68 to 122 of SEQ ID NO:23. The protein having the amino acid sequence shown in SEQ ID NO:20 contains a zinc-finger motif.

Allelic variants of the disclosed subgenomic polynucleotides can occur and encode proteins which are identical, homologous, or substantially related to amino acid sequences disclosed herein (see below).

Allelic variants of subgenomic polynucleotides of the invention can be identified by hybridization of putative allelic variants with nucleotide sequences disclosed herein under stringent conditions. For example, by using the following wash conditions--2 x SCC, 0.1% SDS, room temperature twice, 30 minutes each; then 2 x SCC, 0.1% SDS, 50 °C. once, 30 minutes; then 2 x SCC, room temperature twice, 10 minutes each--allelic variants can be identified which contain at most about 25-30% basepair mismatches. More preferably, allelic variants contain 15-25% basepair mismatches, even more preferably 5-15% basepair mismatches.

Protein variants of secreted proteins of the invention are also included. Amino acids which are not involved in regions which determine biological activity can be deleted or modified without affecting biological function. Preferably, protein

variants of the invention have amino acid sequences which are at least 85%, 90%, or 95% identical to the amino acid sequences disclosed herein and have similar biological properties (see below). More preferably, the molecules are 98% identical. Modifications of interest in the protein sequences can include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue. Proteins or derivatives can be either glycosylated or unglycosylated. Techniques for making such modifications are well known to those skilled in the art (see, e.g., U.S. 4,518,584). Alternatively, variants of proteins disclosed herein can be constructed using techniques of synthetic chemistry or using recombinant DNA methods.

Preferably, amino acid changes in variants or derivatives of proteins of the invention are conservative amino acid changes, *i.e.*, substitutions of similarly charged or uncharged amino acids. A conservative amino acid change involves substitution of one amino acid for another amino acid of a family of amino acids which are structurally related in their side chains. Naturally occurring amino acids are generally divided into four families: acidic (aspartate, glutamate), basic (lysine, arginine, histidine), non-polar (alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), and uncharged polar (glycine, asparagine, glutamine, cystine, serine, threonine, tyrosine) amino acids. Phenylalanine, tryptophan, and tyrosine are sometimes classified as aromatic amino acids. It is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the binding properties of the resulting molecule, especially if the replacement does not involve an amino acid at a binding site involved in an interaction of the protein. Non-naturally occurring amino acids can also be used to form protein variants of the invention.

Whether an amino acid change results in a functional protein or polypeptide can readily be determined by assaying biological properties of the disclosed proteins or polypeptides, as described below. Species homologs of human subgenomic polynucleotides and proteins of the invention can also be identified by making

suitable probes or primers and screening cDNA expression libraries from other species, such as mice, monkeys, yeast, or bacteria.

In the case of proteins which are membrane-bound, such as cell surface receptor proteins, soluble forms of the proteins can be obtained by deleting the nucleotide sequences which encode part or all of the intracellular and transmembrane domains of the protein and expressing a fully secreted form of the protein in a host cell. Techniques for identifying intracellular and transmembrane domains, such as homology searches, can be used to identify such domains in proteins of the invention using amino acid and nucleotide sequences disclosed herein.

Polypeptides consisting of less than full-length proteins of the present invention are also provided. Polypeptides of the invention can be linear or can be cyclized, for example, as described in Saragovi *et al.*, 1992, *Bio/Technology* 10, 773-778 and McDowell *et al.*, 1992, *J. Amer. Chem. Soc.* 114, 9245-9253. Polypeptides can be used, for example, as immunogens, diagnostic aids, or therapeutics, and to create fusion proteins, as described below.

Polypeptide molecules consisting of less than the entire amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38 are also provided. Such polypeptides comprise at least 6, 8, 10, 12, 15, 18, or 20 contiguous amino acids of an amino acid sequence shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38. Polypeptide molecules of the invention can also possess minor amino acid alterations which do not substantially affect the ability of the polypeptides to interact with specific molecules, such as antibodies.

Derivatives of the polypeptides, such as glycosylated forms, aggregative conjugates with other molecules, and covalent conjugates with unrelated chemical moieties, are also provided. Derivatives also include allelic variants, species variants, and muteins. Covalent derivatives are prepared by linkage of functionalities to groups which are found in the amino acid chain or at the N- or C-terminal residue by means known in the art. Truncations or deletions of regions which do not affect biological function are also encompassed. Truncated or deleted

polypeptides can be prepared synthetically or recombinantly, or by proteolytic digestion of purified or partially purified secreted proteins of the invention.

Fusion proteins comprising at least 6, 8, 10, 12, 15, 18, or 20 contiguous amino acids of the disclosed proteins can also be constructed. Human fusion proteins are useful, *inter alia*, for generating antibodies against amino acid sequences and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with secreted proteins of the invention and influence their function. Physical methods, such as protein affinity chromatography, or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and can also be used as drug screens. Fusion proteins can also be used to target molecules to a specific location in a cell or to cause a molecule to be secreted or to be anchored in a cellular membrane.

Fusion proteins of the invention comprise two protein segments which are fused together with a peptide bond. The first protein segment comprises at least 6, 8, 10, 12, 15, 18, or 20 contiguous amino acids selected from an amino acid sequence shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38. The first protein segment can also be a full-length protein (comprising a signal sequence) or a mature protein (lacking a signal sequence). The second protein segment can be a full-length protein or a protein fragment. The second protein or protein fragment can be labeled with a detectable marker, such as a radioactive, chemiluminescent, biotinylated, or fluorescent tag, or can be an enzyme which will generate a detectable product. Enzymes suitable for this purpose, such as  $\beta$ -galactosidase, are well known in the art.

Techniques for making fusion proteins, either recombinantly or by covalently linking two protein segments, are well known in the art. Fusion proteins comprising amino acid sequences of the invention can also be constructed, for example, using standard recombinant DNA methods to make a DNA construct which comprises contiguous nucleotides selected from SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19 and encoding the desired amino

acids in proper reading frame with nucleotides encoding the second protein segment.

Proteins or polypeptides of the invention can be purified free from other components with which they are normally associated in a cell, such as carbohydrates, lipids, subcellular organelles, or other proteins. An isolated protein or polypeptide is at least 90% pure. Preferably, the preparations are 95% or 99% pure. The purity of a preparation can be assessed, for example, by examining electrophoretograms of protein or polypeptide preparations at several pH values and at several polyacrylamide concentrations, as is known in the art.

Standard biochemical methods can be used to isolate proteins of the invention from tissues which express the proteins or to isolate proteins, polypeptides, or fusion proteins from recombinant host cells into which a DNA construct has been introduced. Methods of protein purification, such as size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, crystallization, electrofocusing, or preparative gel electrophoresis, are well known and widely used in the art.

Alternatively, proteins, fusion proteins, or polypeptides of the invention can be produced by recombinant DNA methods or by synthetic chemical methods. Synthetic chemistry methods, such as solid phase peptide synthesis, can be used to synthesize proteins, fusion proteins, or polypeptides. For production of recombinant proteins, fusion proteins, or polypeptides, coding sequences selected from the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19 can be expressed in prokaryotic or eukaryotic host cells using expression systems known in the art. These expression systems include bacterial, yeast, insect, and mammalian cells (see below).

The resulting expressed protein can then be purified from the culture medium or from extracts of the cultured cells using purification procedures known in the art. For example, for proteins fully secreted into the culture medium, cell-free medium can be diluted with sodium acetate and contacted with a cation exchange resin, followed by hydrophobic interaction chromatography. Using this method, the desired protein, fusion protein, or polypeptide is typically greater than 95% pure.



Further purification can be undertaken, using, for example, any of the techniques listed above. Proteins, fusion proteins, or polypeptides can also be tagged with an epitope, such as a "Flag" epitope (Kodak), and purified using an antibody which specifically binds to that epitope.

5                   It may be necessary to modify a protein produced in yeast or bacteria, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain a functional protein. Such covalent attachments can be made using known chemical or enzymatic methods.

10                   Proteins or polypeptides of the invention can also be expressed in cultured cells in a form which will facilitate purification. For example, a secreted protein or polypeptide can be expressed as a fusion protein comprising, for example, maltose binding protein, glutathione-S-transferase, or thioredoxin, and purified using a commercially available kit. Kits for expression and purification of such fusion proteins are available from companies such as New England BioLabs, Pharmacia, and Invitrogen.

15                   The coding sequences disclosed herein can also be used to construct transgenic animals, such as cows, goats, pigs, or sheep. Female transgenic animals can then produce proteins, polypeptides, or fusion proteins of the invention in their milk. Methods for constructing such animals are known and widely used in the art.

20                   Isolated proteins, polypeptides, or fusion proteins of the invention can be used to obtain a preparation of antibodies which specifically bind to epitopes comprising amino acid sequences of the invention. Antibodies of the invention can be used, for example, to detect proteins, polypeptides, or fusion proteins of the invention which are secreted into culture medium or to identify tissues or cells  
25                   which express these molecules. The antibodies can be polyclonal or monoclonal or can be single chain antibodies. Techniques for raising polyclonal and monoclonal antibodies and for constructing single chain antibodies are well known in the art.

30                   Antibodies of the invention bind specifically to epitopes comprising amino acid sequences of the invention, preferably to epitopes not present on other proteins. Typically a minimum number of contiguous amino acids to encode an epitope is 6, 8, or 10. However, more amino acids can be part of an epitope, for

example, at least 15, 25, or 50, especially to form epitopes which involve non-contiguous residues. Specific binding antibodies do not detect other proteins on Western blots of proteins or in immunocytochemical assays. Specific binding antibodies provide a signal at least ten-fold lower than the signal provided with epitopes which do not comprise amino acid sequences of the invention. Antibodies which bind specifically to secreted proteins of the invention include those that bind to mature or full-length proteins, to polypeptides or degradation products, to fusion proteins, or to protein variants. In a preferred embodiment of the invention, the antibodies immunoprecipitate the desired protein, fusion protein, or polypeptide from solution and react with the protein, fusion protein, or polypeptide on Western blots of polyacrylamide gels.

Techniques for purifying antibodies are those which are available in the art. In a preferred embodiment, antibodies are affinity purified by passing the antibodies over a column to which amino acid sequences of the invention are bound. The bound antibody is then eluted, for example using a buffer with a high salt concentration. Any such technique may be chosen to purify antibodies of the invention.

The invention also provides DNA constructs, for expressing all or a portion of a protein of the invention in a host cell. The DNA construct comprises a promoter which is functional in the particular host cell selected. The skilled artisan can readily select an appropriate promoter from the large number of cell type-specific promoters known and used in the art. The DNA construct can also contain a transcription terminator which is functional in the host cell.

The expression construct comprises a polynucleotide segment which encodes all or a portion of a human protein encoded by SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19 or a variant thereof. The polynucleotide segment is located downstream from the promoter. Transcription of the polynucleotide segment initiates at the promoter. DNA constructs can be linear or circular and can contain sequences, if desired, for autonomous replication.

The host cell comprising the DNA construct can be any suitable prokaryotic or eukaryotic cell. Expression systems in bacteria include those described in Chang

*et al.*, *Nature* (1978) 275: 615; Goeddel *et al.*, *Nature* (1979) 281: 544; Goeddel *et al.*, *Nucleic Acids Res.* (1980) 8: 4057; EP 36,776; U.S. 4,551,433; deBoer *et al.*, *Proc. Natl. Acad. Sci. USA* (1983) 80: 21-25; and Siebenlist *et al.*, *Cell* (1980) 20: 269.

5 Expression systems in yeast include those described in Hinnen *et al.*, *Proc. Natl. Acad. Sci. USA* (1978) 75: 1929; Ito *et al.*, *J. Bacteriol.* (1983) 153: 163; Kurtz *et al.*, *Mol. Cell. Biol.* (1986) 6: 142; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25: 141; Gleeson *et al.*, *J. Gen. Microbiol.* (1986) 132: 3459, Roggenkamp *et al.*, *Mol. Gen. Genet.* (1986) 202 :302); Das *et al.*, *J. Bacteriol.* (1984) 158: 1165; De Louvencourt *et al.*, *J. Bacteriol.* (1983) 154: 737, Van den Berg *et al.*, *Bio/Technology* (1990) 8: 135; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25: 141; Cregg *et al.*, *Mol. Cell. Biol.* (1985) 5: 3376; U.S. 4,837,148; U.S. 4,929,555; Beach and Nurse, *Nature* (1981) 300: 706; Davidow *et al.*, *Curr. Genet.* (1985) 10: 380; Gaillardin *et al.*, *Curr. Genet.* (1985) 10: 49; Ballance *et al.*, *Biochem. Biophys. Res. Commun.* (1983) 112: 284-289; Tilburn *et al.*, *Gene* (1983) 26: 205-22,, Yelton *et al.*, *Proc. Natl. Acad. Sci. USA* (1984) 81: 1470-1474; Kelly and Hynes, *EMBO J.* (1985) 4: 475479; EP 244,234; and WO 91/00357.

20 Expression of heterologous genes in insects can be accomplished as described in U.S. 4,745,051; Friesen *et al.* (1986) "The Regulation of Baculovirus Gene Expression" in: THE MOLECULAR BIOLOGY OF BACULOVIRUSES (W. Doerfler, ed.); EP 127,839; EP 155,476; Vlak *et al.*, *J. Gen. Virol.* (1988) 69: 765-776; Miller *et al.*, *Ann. Rev. Microbiol.* (1988) 42: 177; Carbonell *et al.*, *Gene* (1988) 73: 409; Maeda *et al.*, *Nature* (1985) 315: 592-594; Lebacq-Verheyden *et al.*, *Mol. Cell. Biol.* (1988) 8: 3129; Smith *et al.*, *Proc. Natl. Acad. Sci. USA* (1985) 82: 8404; Miyajima *et al.*, *Gene* (1987) 58: 273; and Martin *et al.*, *DNA* (1988) 7:99. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts are described in Luckow *et al.*, *Bio/Technology* (1988) 6: 47-55, Miller *et al.*, in GENERIC ENGINEERING (Setlow, J.K. *et al.* eds.), Vol. 8 (Plenum Publishing, 1986), pp. 277-279; and Maeda *et al.*, *Nature*, (1985) 315: 592-594.

30 Mammalian expression can be accomplished as described in Dijkema *et al.*,

EMBO J. (1985) 4: 761; Gorman *et al.*, *Proc. Natl. Acad. Sci. USA* (1982b) 79: 6777; Boshart *et al.*, *Cell* (1985) 41: 521; and U.S. 4,399,216. Other features of mammalian expression can be facilitated as described in Ham and Wallace, *Meth. Enz.* (1979) 58: 44; Barnes and Sato, *Anal. Biochem.* (1980) 102: 255; U.S. 4,767,704; U.S. 4,657,866; U.S. 4,927,762; U.S. 4,560,655; WO 90/103430, WO 87/00195, and U.S. RE 30,985.

DNA constructs of the invention can be introduced into host cells using any technique known in the art. These techniques include transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, and calcium phosphate-mediated transfection.

Alternatively, expression of an endogenous gene encoding a protein of the invention can be manipulated by introducing by homologous recombination a DNA construct comprising a transcription unit in frame with the endogenous gene, to form a homologously recombinant cell comprising the transcription unit. The transcription unit comprises a targeting sequence, a regulatory sequence, an exon, and an unpaired splice donor site. The new transcription unit can be used to turn the endogenous gene on or off as desired. This method of affecting endogenous gene expression is taught in U.S. 5,641,670, which is incorporated herein by reference.

The targeting sequence is a segment of at least 10, 12, 15, 20, or 50 contiguous nucleotides selected from the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19. The transcription unit is located upstream to a coding sequence of the endogenous gene. The exogenous regulatory sequence directs transcription of the coding sequence of the endogenous gene.

Secreted proteins of the invention have a variety of uses. For example, secreted proteins can be used in assays to determine biological activities, such as cytokine, cell proliferation, or cellular differentiation activities, tissue growth or

regeneration, activin or inhibin activity, chemotactic or chemokinetic activity, hemostatic or thrombolytic activity, receptor/ligand activity, tumor inhibition, or anti-inflammatory activity. Assays for these activities are known in the art and are disclosed, for example, in U.S. 5,654,173, which is incorporated herein by reference.

Proteins of the invention can also be used as biomarkers, to identify tissues or cell types which express the proteins, or a stage- or disease-specific alteration in protein expression. Proteins of the invention can be used in protein interaction assays, to identify ligands or binding proteins. Compounds which affect the biological activities of the secreted proteins or their ability to interact with specific ligands can be identified using proteins of the invention in screening assays. Proteins and antibodies of the invention can also be used to design diagnostic tests and therapeutic compositions for diseases which may be associated with altered expression of these proteins. Fusion proteins comprising, for example, signal sequences or transmembrane domains of the disclosed proteins, can be used to target other protein domains to cellular locations in which the domains are not normally found, such as bound to a cellular membrane or secreted extracellularly.

Further objects, features, and advantages of the present invention will readily occur to the skilled artisan provided with the disclosure above.

## **SYNOPSIS OF THE INVENTION**

1. An isolated and purified human protein having an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

2. An isolated and purified human protein having an amino acid sequence which is at least 85% identical to an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

3. The isolated and purified human protein of item 2 wherein the amino acid sequence is at least 90% identical.

4. The isolated and purified human protein of item 2 wherein the amino acid sequence is at least 95% identical.

5. The isolated and purified human protein of item 2 wherein the amino acid sequence is at least 98% identical.

6. An isolated and purified human polypeptide comprising at least 6 contiguous amino acids of an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

7. A fusion protein comprising a first protein segment and a second protein segment fused together by means of a peptide bond, wherein the first protein segment consists of at least 6 contiguous amino acids selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38.

8. A preparation of antibodies which specifically bind to the human protein of item 1.

9. The preparation of antibodies of item 8 wherein the antibodies are monoclonal.

10. The preparation of antibodies of item 8 wherein the antibodies are polyclonal.

11. The preparation of antibodies of item 8 wherein the antibodies are single chain antibodies.

12. An isolated and purified subgenomic polynucleotide having a nucleotide sequence selected from the group consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

13. An isolated and purified subgenomic polynucleotide consisting of at least 10 contiguous nucleotides of a nucleotide sequence selected from the group

consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

14. An isolated gene corresponding to a cDNA sequence selected from the group consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19.

15. A DNA construct for expressing all or a portion of a human protein having an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38, comprising:

a promoter; and

a polynucleotide segment encoding at least 6 contiguous amino acids of the human protein, wherein the polynucleotide segment is located downstream from the promoter, wherein transcription of the polynucleotide segment initiates at or 3' to the promoter.

16. A host cell comprising a DNA construct comprising:

a promoter; and

a polynucleotide segment encoding at least 6 contiguous amino acids of a human protein having an amino acid sequence selected from the group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38, wherein the polynucleotide segment is located downstream from the promoter and wherein transcription of the polynucleotide segment initiates at or 3' to the promoter.

17. A homologously recombinant cell having incorporated therein a new transcription initiation unit, wherein the new transcription initiation unit comprises in 5' to 3' order:

(a) an exogenous regulatory sequence;

(b) an exogenous exon; and

(c) a splice donor site,

wherein the transcription initiation unit is located upstream to a coding sequence of a gene, wherein the gene comprises a nucleotide sequence selected from the group

consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19, and wherein the exogenous regulatory sequence controls transcription of the coding sequence of the gene.

5 18. A method of producing a human protein, comprising the steps of:  
growing a culture of a cell comprising a DNA construct comprising  
(1) a promoter and (2) a polynucleotide segment encoding at least 6 contiguous  
amino acids of a human protein having an amino acid sequence selected from the  
group consisting of the amino acid sequences shown in SEQ ID Nos:20, 21, 22, 23,  
24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, and 38, wherein the  
10 polynucleotide segment is located downstream from the promoter and wherein  
transcription of the polynucleotide segment initiates at or 3' to the promoter; and;  
purifying the protein from the culture.

15 19. A method of producing a human protein, comprising the steps of:  
growing a culture of a homologously recombinant cell having  
incorporated therein a new transcription initiation unit, wherein the new  
transcription initiation unit comprises in 5' to 3' order:

- (a) an exogenous regulatory sequence;
- (b) an exogenous exon; and
- (c) a splice donor site,

20 wherein the transcription initiation unit is located upstream to a coding sequence of  
a gene, wherein the gene comprises a nucleotide sequence selected from the group  
consisting of the nucleotide sequences shown in SEQ ID NOs:1, 2, 3, 4, 5, 6, 7, 8,  
9, 10, 11, 12, 13, 14, 15, 16, 17, 18, and 19 and wherein the exogenous regulatory  
sequence controls transcription of the coding sequence of the gene; and  
25 purifying the protein from the culture.

20. A method of identifying a secreted polypeptide which is modified by  
rough microsomes, comprising the steps of:  
transcribing *in vitro* a population of cDNA molecules whereby a  
population of cRNA molecules is formed;



translating a first portion of the population of cRNA molecules *in vitro* in the absence of rough microsomes whereby a first population of polypeptides is formed;

5 translating a second portion of the population of cRNA molecules *in vitro* in the presence of rough microsomes whereby a second population of polypeptides is formed;

comparing the first population of polypeptides with the second population of polypeptides; and

10 detecting polypeptide members of the second population which have been modified by the rough microsomes.

21. The method of item 20 wherein the population of cDNA molecules is synthesized by reverse transcription of a population of mRNA molecules.

22. The method of item 21 wherein the mRNA molecules are isolated from a mammal.

15 23. The method of item 22 wherein the mRNA molecules are isolated from a human.

24. The method of item 20 wherein the population of cDNA molecules is obtained from a cDNA library.

20 25. The method of item 24 wherein the cDNA library is derived from a mammalian genome.

26. The method of item 25 wherein the cDNA library is derived from a human genome.

## SEQUENCE LISTING

### (1) GENERAL INFORMATION

- (i) APPLICANT: Escobedo, Jaime  
Quianjin, Hu  
Garcia, Pablo  
Williams, Lewis T.  
Kothakota, Srinivas
- (ii) TITLE OF THE INVENTION: Secreted Human Proteins
- (iii) NUMBER OF SEQUENCES: 38
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Banner & Witcoff
  - (B) STREET: 1001 G Street, NW
  - (C) CITY: Washington
  - (D) STATE: DC
  - (E) COUNTRY: USA
  - (F) ZIP: 20001
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Diskette
  - (B) COMPUTER: IBM Compatible
  - (C) OPERATING SYSTEM: DOS
  - (D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE: 11-DEC-1997

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 60/032757

(B) FILING DATE: 11-DEC-1996

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Kagan, Sarah A

(B) REGISTRATION NUMBER: 32141

(C) REFERENCE/DOCKET NUMBER:

2441.39505;1369.002;1452.001

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 202-508-9100

(B) TELEFAX: 202-508-9299

(C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2063 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAATTCGGCA	CGAGGCCTCA	GTCTTCCAGG	GCGGCGGTGG	GTGTCCGCTT	CTCTCTGCTC	60
TTCGACTGCA	CCGCACTCGC	GCGTGACCTT	GACTCCCCCT	AGTCAGCTCA	GCGGTGCTGC	120
CATGGCGTGG	CGGCGGCGCG	AAGCCGGCGT	CGGGGCTCGC	GGCGTGTTGG	CTCTGGCGTT	180
GCTCGCCCTG	GCCCTGTGCG	TGCCCGGGGC	COGGGGCCGG	GCTCTCGAGT	GGTTCTCGGC	240

CGTGGTAAAC	ATCGAGTACG	TGGACCCGCA	GACCAACCTG	ACGGTGTGGA	GCGTCTCGGA	300
GAGTGGCCGC	TTCGGCGACA	GCTCGCCCAA	GGAGGGCGCG	CATGGCCTGG	TGGGCGTCCC	360
GTGGGCGCCC	GGCGGAGACC	TCGAGGGCTG	CGCGCCCGAC	ACGCGCTTCT	TCGTGCCCGA	420
GCCCCGCGGC	CGAGGGGCGG	CGCCCTGGGT	CGCCCTGGTG	GCTCGTGGGG	GCTGCACCTT	480
CAAGGACAAG	GTGCTGGTGG	CGGCGCGGAG	GAACGCCTCG	GCCGTCGTCC	TCTACAATGA	540
GGAGCGCTAC	GGGAACATCA	CCTTGCCCAT	GTCTCACGCG	GGAACAGGAA	ATATAGTGGT	600
CATTATGATT	AGCTATCCAA	AAGGAAGAGA	AATTTTGGAG	CTGGTGCAA	AAGGAATTCC	660
AGTAACGATG	ACCATAGGGG	TTGGCACCOG	GCATGTACAG	GAGTTCATCA	GCGGTTCAGTC	720
TGTGGTGTTT	GTGGCCATTG	CCTTCATCAC	CATGATGATT	ATCTCGTTAG	CCTGGCTAAT	780
ATTTTACTAT	ATACAGCGTT	TCCTATATAC	TGGCTCTCAG	ATTGGAAGTC	AGAGCCATAG	840
AAAAGAACT	AAGAAAGTTA	TTGGCCAGCT	TCTACTTCAT	ACTGTAAAGC	ATGGAGAAAA	900
GGGAATTGAT	GTTGATGCTG	AAAATTGTGC	AGTGTGTATT	GAAAATTTCA	AAGTAAAGGA	960
TATTATTAGA	ATTCTGCCAT	GCAAGCATAT	TTTTCATAGA	ATATGCATTG	ACCCATGGCT	1020
TTTGGATCAC	CGAACATGTC	CAATGTGTAA	ACTTGATGTC	ATCAAAGCCC	TAGGATATTG	1080
GGGAGAGCCT	GGGGATGTAC	AGGAGATGCC	TGCTCCAGAA	TCTCCTCCTG	GAAGGGATCC	1140
AGCTGCAAAT	TTGAGTCTAG	CTTTACCAGA	TGATGACGGA	AGTGATGACA	GCAGTCCACC	1200
ATCAGCCTCC	CCTGCTGAAT	CTGAGCCACA	GTGTGATCCC	AGCTTTAAAG	GAGATGCAGG	1260
AGAAAATACG	GCATTGCTAG	AAGCCGGCAG	GAGTGACTCT	CGGCATGGAG	GACCCATCTC	1320
CTAGCACACG	TGCCCCTGA	AGTGGCACCA	ACAGAAGTTT	GGCTTGAAC	AAAGGACATT	1380
TTATTTTTTT	TACTTTAGCA	CATAATTTGT	ATATTTGAAA	ATAATGTATA	TTATTTTACC	1440
TATTAGATTC	TGATTTGATA	TACAAAGGAC	TAAGATATTT	TCTTCTTGAA	GAGACTTTTC	1500
GATTAGTCCT	CATATATTTA	TCTACTAAAA	TAGAGTGTTC	ACCATGAACA	GTGTGTTGCT	1560
TCAGACTATT	ACAAAGACAA	CTGGGGCAGG	TACTCTAATA	TAAAGGACAG	GTGGTGTTTC	1620
TAAATAATTG	GCTGCTATGG	TTCTGTAAAA	ACCAGTTAAT	TCTATTTTTC	AAGGTTTTTG	1680
GCAAAGCACA	TCAATGTTAG	ACTAGTTGAA	GTGGAATTGT	ATAATTCAAT	TCGATAATTG	1740
ATCTCATGGG	CTTTCCCTGG	AGGAAAGGTT	TTTTTTGTTG	TTTTTTTTTT	AAGAACTTGA	1800
AACCTGTAAA	CTGAGATGTC	TGTAGCTTTT	TTGCCCATCT	GTAGTGTATG	TGAAGATTTC	1860
AAAACCTGAG	AGCACTTTTT	CTTTGTTTGT	AATTATGAGA	AAGGCACTAG	ATGACTTTAG	1920
GATTTGCATT	TTTCCCTTTA	TTGCCTCATT	TCTTGTGACG	CCTTGTGTTG	GAGGGAAATC	1980
TGTTTATTTT	TTCTACAAA	TAAAAAGCTA	AGATTCTATA	TCGCAAAAAA	AAAAAAAAAA	2040
AAAAAAAAAA	TTCCTGCGGC	CGC				2063

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1328 base pairs

- (B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GAATTTCGGCA	CGAGGTAGGC	AAGGGATAAA	AAGGCACCTA	AGGCCCTTTT	GCAATAAGAA	60
GCCAGATGGA	TAAAGGAAGT	GCTGGTCACC	CTGGAGGTGT	ACTGGTTTGG	GGAAGGTCCC	120
CGGCCCCCAC	AGCCCTCTGG	GGAGCCTCAC	CCTGGCTCTC	CCCACTCACC	TCAGCCCTCA	180
GGCAGCCCCT	CCACAGGGCC	CCTCTCCTGC	CTGGACAGCT	CTGCTGGTCT	CCCCGTCCCC	240
TGGAGAAGAA	CAAGGCCATG	GGTCGGCCCC	TGCTGCTGCC	CCTGCTGCTC	CTGCTGCAGC	300
CGCCAGCATT	TCTGCAGCCT	GGTGGCTCCA	CAGGATCTGG	TCCAAGCTAC	CTTTATGGGG	360
TCACTCAACC	AAAACACCTC	TCAGCCTCCA	TGGGTGGCTC	TGTGGAAATC	CCCTTCTCCT	420
TCTATTACCC	CTGGGAGTTA	GCCATAGTTC	CCAACGTGAG	AATATCCTGG	AGACGGGGCC	480
ACTTCCACGG	GCAGTCCTTC	TACAGCACAA	GGCCGCCTTC	CATTCAACAAG	GATTATGTGA	540
ACCGGCTCTT	TCTGAACTGG	ACAGAGGGTC	AGGAGAGCGG	CTTCCTCAGG	ATCTCAAACC	600
TGCGGAAGGA	GGACCAGTCT	GTGTATTTCT	GCCGAGTCGA	GCTGGACACC	CGGAGATCAG	660
GGAGGCAGCA	GTTGCAGTCC	ATCAAGGGGA	CCAACTCAC	CATCACCAG	GCTGTCACAA	720
CCACCACCAC	CTGGAGGCC	AGCAGCACAA	CCACCATAGC	CGGCCTCAGG	GTCACAGAAA	780
GCAAAGGGCA	CTCAGAATCA	TGGCACCTAA	GTCTGGACAC	TGCCATCAGG	GTTGCATTGG	840
CTGTGCTGT	GCTCAAACT	GTCATTTTGG	GACTGCTGTG	CCTCCTCCTC	CTGTGGTGGA	900
GGAGAAGGAA	AGGTAGCAGG	GCGCCAAGCA	GTGACTTCTG	ACCAACAGAG	TGTGGGGAGA	960
AGGGATGTGT	ATTAGCCCCG	GAGGACGTGA	TGTGAGACCC	GCTTGTGAGT	CCTCCACACT	1020
CGTTCCCCAT	TGGCAAGATA	CATGGAGAGC	ACCCTGAGGA	CCTTTAAAAG	GCAAAGCCGC	1080
AAGGCAGAAG	GAGGCTGGGT	CCCTGAATCA	CCGACTGGAG	GAGAGTTACC	TACAAGAGCC	1140
TTCATCCAGG	AGCATCCACA	CTGCAATGAT	ATAGGAATGA	GGTCTGAACT	CCACTGAATT	1200
AAACCACTGG	CATTGTTTTG	CTGTTTATTA	TAGCAGTGCA	AAGAGTTCCT	TTATCCTCCC	1260
CAAGGATGGA	AAAATACAAT	TTATTTTGCT	TACCATAAAA	AAAAAAAAAA	AAAAATTCCT	1320
GCGGCCGC						1328

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1689 base pairs  
(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAATTCGGCA	CGAGGGCAAG	ATTCGATACA	AAACCAATGA	ACCTGTGTGG	GAGGAAACT	60
TCACTTTCTT	CATTACAAT	CCCAAGCGCC	AGGACCTTGA	AGTTGAGGTC	AGAGACGAGC	120
AGCACCAGTG	TTCCCTGGGG	AACCTGAAGG	TCCCCCTCAG	CCAGCTGCTC	ACCAAGTGAGG	180
ACATGACTGT	GAGCCAGCGC	TTCCAGCTCA	GTAACCTCGG	TCCAAACAGC	ACCATCAAGA	240
TGAAGATTGC	CCTGCGGGTG	CTCCATCTCG	AAAAGCGAGA	AAGGCCTCCA	GACCACCAAC	300
ACTCAGCTCA	AGTCAAACGT	CCCTCTGTGT	CCAAAGAGGG	GAGGAAAACA	TCCATCAAAT	360
CTCATATGTC	TGGGTCTCCA	GGCCCTGGTG	GCAGCAACAC	AGCTCCATCC	ACACCAGTCA	420
TTGGGGGCGC	TGATAAGCCT	GGTATGGAAG	AAAAGGCCCA	GCCCCCTGAG	GCCGGCCCTC	480
AGGGGCTGCA	CGACCTGGGC	AGAAGCTCCT	CCAGCCTCCT	GGCCTCCCCA	GGCCACATCT	540
CAGTCAAGGA	GCCGACCCCC	AGCATCGCCT	CGGACATCTC	GCTGCCCATC	GCCACCCAGG	600
AGCTGCGGCA	AAGGCTGAGG	CAGCTGGAAA	ACGGGACGAC	CCTGGGACAG	TCTCCACTGG	660
GGCAGATCCA	GCTGACCATC	CGGCACAGCT	CGCAGAGAAA	CAAGCTTATC	GTGGTCTGTC	720
ATGCCCTGCAG	AAACCTCATT	GCCTTCTCTG	AAGACGGCTC	TGACCCCTAT	GTCCGCATGT	780
ATTTATTACC	AGACAAGAGG	CGGTCAGGAA	GGAGGAAAAC	ACACGTGTCA	AAGAAAACAT	840
TAAATCCAGT	GTTTGATCAA	AGCTTTGATT	TCAGTGTTC	GTTACCAGAA	GTGCAGAGGA	900
GAACGCTCGA	CGTTGCCGTG	AAGAACAGTG	GCGGCTTCCT	GTCCAAAGAC	AAAGGGCTCC	960
TTGGCAAAGT	ATTGGTTGCT	CTGGCATCTG	AAGAACTTGC	CAAAGGCTGG	ACCCAGTGGT	1020
ATGACCTCAC	GGAAGATGGG	ACGAGGCCTC	AGGCGATGAC	ATAGCCGCAG	CAGGCAGGAG	1080
GCGTCCTCTT	CAGCGTAGCT	CTCCACCTCT	ACCCGGAACA	CACCCTCTCA	CAGACGTACC	1140
AATGTTATTT	TTATAATTTT	ATGGATTTAG	TTATACATAC	CTTAATAGTT	TTATAAAATT	1200
GTTGACATTT	CAGGCAAATT	TGGCCAATAT	TATCATTGAA	TTTTCTGTGT	TGGATTTTCT	1260
CTAGGATTTT	GCCAGTTCCT	ACAACGTGCA	GTAGGGCGGC	GGTAGCTCTT	GTGTCTGTGG	1320
ACTCTGCTCA	GCTGTGTCCG	TAGGAGTCGG	ATGTGTCTGT	GCTTTATTAT	GGCCTTGTTT	1380
ATATATCACT	GAGGTATACT	ATGCCATGTA	AATAGACTAT	TTTTTATAAT	CTTAACATGC	1440
TGGTTTAAAT	TCAGAAGGAA	ATAGATCAAG	GAAATATATA	TATTTTCTTC	TAAAACTTAT	1500
TAAATTCGTG	TGACAAATAA	TCATTTTCAT	CTTGGCAGCA	AAAAGTTCTC	AGTGACCTAT	1560
TTTGTGGTGT	TTCTTTTTGA	AAAGAAAAGC	TGAAATATTA	TTAAATGCTA	GTATGTTTCT	1620
GCCCATATAT	AAAGATGAAA	TAAAGTATTC	AAAATATTAA	AAAAAAAAAA	AAAAAATTC	1680
TGCGGCCCGC						1689

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1505 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GAATTCGGCA	CGAGGAGCAG	ATCTGCAAGA	GTTTCGTTTA	TGGAGGCTGC	TTGGGCAACA	60
AGAACAATA	CCTTCGGGAA	GAAGAGTGCA	TTCTAGCCTG	TCGGGGTGTG	CAAGGTGGGC	120
CTTTGAGAGG	CAGCTCTGGG	GCTCAGGCGA	CTTTCCCCCA	GGGCCCCTCC	ATGGAAAGGC	180
GCCATCCAGT	GTGCTCTGGC	ACCTGTCAGC	CCACCCAGTT	CCGCTGCAGC	AATGGCTGCT	240
GCATCGACAG	TTTCTGGAG	TGTGACGACA	CCCCCAACTG	CCCCGACGCC	TCCGACGAGG	300
CTGCCTGTGA	AAAATACACG	AGTGGCTTTG	ACGAGCTCCA	GCGCATCCAT	TTCCCCAGCG	360
ACAAAGGGCA	CTGCGTGGAC	CTGCCAGACA	CAGGACTCTG	CAAGGAGAGC	ATCCCGCGCT	420
GGTACTACAA	CCCCTTCAGC	GAACACTGCG	CCCCTTTTAC	CTATGGTGGT	TGTTACGGCA	480
ACAAGAACAA	CTTTGAGGAA	GAGCAGCAGT	GCCTCGAGTC	TTGTCGCGGC	ATCTCCAAGA	540
AGGATGTGTT	TGGCCTGAGG	CGGGAAATCC	CCATTCCCAG	CACAGGCTCT	GTGGAGATGG	600
CTGTGCGAGT	GTTCTGGTGC	ATCTGCATTG	TGGTGGTGGT	AGCCATCTTG	GGTTACTGCT	660
TCTTCAAGAA	CCAGAGAAAG	GACTTCCACG	GACACCACCA	CCACCACCA	CCCACCCCTG	720
CCAGCTCCAC	TGTCTCCACT	ACCGAGGACA	CGGAGCACCT	GGTCTATAAC	CACACCACGC	780
GGCCCCCTCTG	AGCCTGGGTC	TCACCGGCTC	TCACCTGGCC	CTGCTTCCTG	CTTGCCAAGG	840
CAGAGGCCCTG	GGCTGGGAAA	AACTTTGGAA	CCAGACTCTT	GCCTGTTTCC	CAGGCCCACT	900
GTGCCTCAGA	GACCAGGGCT	CCAGCCCCTC	TTGGAGAAGT	CTCAGCTAAG	CTCACGTCCT	960
GAGAAAGCTC	AAAGGTTTGG	AAGGAGCAGA	AAACCCTTGG	GCCAGAAGTA	CCAGACTAGA	1020
TGGACCTGCC	TGCATAGGAG	TTTGGAGGAA	GTTGGAGTTT	TGTTTCCTCT	GTTCAAAGCT	1080
GCCTGTCCCT	ACCCCATGGT	GCTAGGAAGA	GGAGTGGGGT	GGTGTGAGAC	CCTGGAGGCC	1140
CCAACCCTGT	CCTCCCGAGC	TCCTCTTCCA	TGCTGTGCGC	CCAGGGCTGG	GAGGAAGGAC	1200
TTCCCTGTGT	AGTTTGTGCT	GTAAAGAGTT	GCTTTTGTGT	TATTTAATGC	TGTGGCATGG	1260
GTGAAGAGGA	GGGGAAGAGG	CCTGTTTGGC	CTCTCTATCC	TCTCTTCTC	TTCCCCCAAG	1320
ATTGAGCTCT	CTGCCCTTGA	TCAGCCCCAC	CCTGGCCTAG	ACCAGCAGAC	AGAGCCAGGA	1380
GAAGCTCAGC	TGCATTCCGC	AGCCCCCACC	CCCAAGGTTT	TCCAACATCA	CAGCCCAGCC	1440
CGCCCCACTGG	GTAATAAAAG	TGGTTTGTGG	AAAAAAAAAA	AAAAAAAAAA	AAGTCCTGCG	1500

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2002 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GAATTCGGCA	CGAGGGCCAT	GGCCGGGCTA	TCCCGCGGGT	CCGCGCGCGC	ACTGCTCGCC	60
GCCCTGCTGG	CGTCGACGCT	GTTGGCGCTG	CTCGTGTCGC	CCGCGCGGGG	TCGCGGCGGC	120
CGGGACCACG	GGGACTGGGA	CGAGGCCTCC	CGGCTGCCGC	CGCTACCACC	CCGCGAGGAC	180
GCGGCGCGCG	TGGCCCGCTT	CGTGACGCAC	GTCTCCGACT	GGGGCGCTCT	GGCCACCATC	240
TCCACGCTGG	AGGCGGTGCG	CGGCCGGCCC	TTCGCCGACG	TCCTCTCGCT	CAGCGACGGG	300
CCCCCGGGCG	CGGGCAGCGG	CGTGCCCTAT	TTCTACCTGA	GCCCGCTGCA	GCTCTCCGTG	360
AGCAACCTGC	AGGAGAATCC	ATATGCTACA	CTGACCATGA	CTTTGGCACA	GACCAACTTC	420
TGCAAGAAAC	ATGGATTTGA	TCCACAAAGT	CCCCTTTGTG	TTCACATAAT	GCTGTCAGGA	480
ACTGTGACCA	AGGTGAATGA	AACAGAAATG	GATATTGCAA	AGCATTGCTT	ATTCATTCTGA	540
CACCCTGAGA	TGAAAACCTG	GCCTTCCAGC	CATAATTGGT	TCTTTGCTAA	GTTGAATATA	600
ACCAATATCT	GGGTCCTGGA	CTACTTTGGT	GGACCAAAAA	TCGTGACACC	AGAAGAATAT	660
TATAATGTCA	CAGTTCAGTG	AAGCAGACTG	TGGTGAATTT	AGCAACACTT	ATGAAGTTTC	720
TTAAAGTGGC	TCATACACAC	TTAAAAGGCT	TAATGTTTCT	CTGGAAAGCG	TCCCAGAATA	780
TTAGCCAGTT	TTCTGTCACA	TGCTGGTTTG	TTTGCTTGCT	TGTTTACTTG	CTTGTTTACC	840
AATAGAGTTG	ACCTGTTATT	GGATTTCTTG	GAAGATGTGG	TAGCTACTTT	TTTCCTATTT	900
TGAAGCCATT	TTCTGTAGAG	AATATCCTTC	ACTATAATCA	AATAAGTTTT	GTCCCATCAA	960
TTCCAAAGAT	GTTTCCAGTG	GTGCTCTTGA	AGAGGAATGA	GTACCAGTTT	TAAATTGCCC	1020
ATTGGCATT	GAAGGTAGTT	GAGTATGTGT	TCTTTATTCC	TAGAAGCCAC	TGTGCTTGGT	1080
AGAGTGCATC	ACTCACCACA	GCTGCCTCTT	GAGCTGCCTG	AGCCTGGTGC	AAAAGGATTG	1140
GCCCCCATTA	TGGTGCTTCT	GAATAAATCT	TGCCAAGATA	GACAAACAAT	GATGAAACTC	1200
AGATGGAGCT	TCCTACTCAT	GTTGATTTAT	GTCTCACAAT	CCTGGGTATT	GTTAATTCAA	1260
CATAGGGTGA	AACTATTTCT	GATAAAGAAC	TTTTGAAAAA	CTTTTTTATAC	TCTAAAGTGA	1320
TACTCAGAAC	AAAAGAAAGT	CATAAACTC	CTGAATTTAA	TTTCCCCACC	TAAGTCGAGA	1380



CAGTATTATC	AAAACACATG	TGCACACAGA	TTATTTTTTG	GCTCCAAAAC	TGGATTGCAA	1440
AAGAAAGAGG	AGAGATATTT	TGTGTGTTCC	TGGTATTCTT	TTATAAGTAA	AGTTACCCAG	1500
GCATGGACCA	GCTTCAGCCA	GGGACAAAAT	CCCCTCCCAA	ACCACTCTCC	ACAGCTTTTT	1560
AAAAATACTT	CTACTCTTAA	CAATTACCTA	AGGTTCCTTC	AAACCCCCCC	AACTCTTAAT	1620
AGCTTCTAGT	GCTGCTACAA	TCTAAGTCAG	GTCACCAGAG	GGAAGAGAAC	ATGGCATTAA	1680
AAGAATCACA	TCTTCAGAAG	AGAAGACACT	AATATTATTA	CCCATATACA	TGATTCAGAG	1740
AGATGACATA	AGATTCCTCT	TAAAGAGGAA	ATGTCAGGAA	TCAAGCCACT	GAATCCCTAA	1800
AGAGAAAAGT	TGAATATGAG	TCATTGTGTC	TGAAAACCTGC	AAAGTGAAC	TAACTGAGAT	1860
CCAGCAAACA	GGTTCGTGTT	AAGAAAAATA	ATTTATACTA	AATTTAGTAA	AATGGACTTC	1920
TTATTCAAAG	CATCAATAAT	TAAAAGAATT	ATTTTAAAAA	AAAAAAAAAA	AAAAAAAAAA	1980
AAAAAAAAAT	TCCTGCGGCC	GC				2002

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1322 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GAATTCGGCA	CGAGGGCCAC	GACTCTGCTG	GCATTTCTTC	TATAGCCACT	GGAATCTGAT	60
CCTGATTGTC	TTCCACTACT	ACCAGGCCAT	CACCACTCCG	CCTGGGTACC	CACCCCAGGG	120
CAGGAATGAT	ATCGCCACCG	TCTCCATCTG	TAAGAAGTGC	ATTTACCCCA	AGCCAGCCCG	180
AACACACCAC	TGCAGCATCT	GCAACAGGTG	TGTGCTGAAG	ATGGATCACC	ACTGCCCTG	240
GCTAAACAAT	TGTGTGGGCC	ACTATAACCA	TCGGTACTTC	TTCTCTTTCT	GCTTTTTCAT	300
GACTCTGGGC	TGTGTCTACT	GCAGCTATGG	AAGTTGGGAC	CTTTTCCGGG	AGGCTTATGC	360
TGCCATTGAG	AAAATGAAAC	AGCTCGACAA	GAACAACTA	CAGGCGGTTG	CCAACCAGAC	420
TTATCACCAG	ACCCACCCAC	CCACCTTCTC	CTTTCGAGAA	AGGATGACTC	ACAAGAGTCT	480
TGTCTACCTC	TGGTTCCTGT	GCAGTTCTGT	GGCACTTGCC	CTGGGTGCCC	TAACTGTATG	540
GCATGCTGTT	CTCATCAGTC	GAGGTGAGAC	TAGCATCGAA	AGGCACATCA	ACAAGAAGGA	600
GAGACGTCGG	CTACAGGCCA	AGGGCAGAGT	ATTTAGGAAT	CCTTACAAC	ACGGCTGCTT	660
GGACAACTGG	AAGGTATTCC	TGGGTGTGGA	TACAGGAAGG	CACTGGCTTA	CTCGGGTGCT	720
CTTACCTTCT	ACTCACTTGC	CCCATGGGAA	TGGAATGAGC	TGGGAGCCCC	CTCCCTGGGT	780

GACTGCTCAC	TCAGCCTCTG	TGATGGCAGT	GTGAGCTGGA	CTGTGTCAGC	CACGACTCGA	840
GCACTCATTC	TGCTCCCTAT	GTTATTTCAA	GGGCCTCCAA	GGGCAGCTTT	TCTCAGAATC	900
CTTGATCAAA	AAGAGCCAGT	GGGCCTGCCT	TAGGGTACCA	TGCAGGACAA	TTCAAGGACC	960
AGCCTTTTTA	CCACTGCAGA	AGAAAGACAC	AATGTGGAGA	AATCTTAGGA	CTGACATCCC	1020
TTTACTCAGG	CAAACAGAAG	TTCCAACCCC	AGACTAGGGG	TCAGGCAGCT	AGCTACCTAC	1080
CTTGCCCAGT	GCTGACCCGG	ACCTCCTCCA	GGATACAGCA	CTGGAGTTGG	CCACCACCTC	1140
TTCTACTTGC	TGTCTGAAAA	AACACCTGAC	TAGTACAGCT	GAGATCTTGG	CTTCTCAACA	1200
GGGCAAAGAT	ACCAGGCCTG	CTGCTGAGGT	CACTGCCACT	TCTCACATGC	TGCTTAAGGG	1260
AGCACAAATA	AAGGTATTCG	ATTTTAAAAA	AAAAAAAAAA	AAAAAAAAAT	TCCTGCGGCC	1320
GC						1322

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1573 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GAATTCGGCA	CGAGGAGCCT	GCCTTCATCT	AGGATGGCTC	CTCTGGGCAT	GCTGCTTGGG	60
CTGCTGATGG	CCGCCTGCTT	CACCTTCTGC	CTCAGTCATC	AGAACCTGAA	GGAGTTTGCC	120
CTGACCAACC	CAGAGAAGAG	CAGCACCAAA	GAAACAGAGA	GAAAAGAAAC	CAAAGCCGAG	180
GAGGAGCTGG	ATGCCGAAGT	CCTGGAGGTG	TTCCACCCGA	CGCATGAGTG	GCAGGCCCTT	240
CAGCCAGGGC	AGGCTGTCCC	TGCAGGATCC	CACGTACGGC	TGAATCTTCA	GACTGGGGAA	300
AGAGAGGCAA	AACTCCAATA	TGAGGACAAG	TTCCGAAATA	ATTGAAAGG	CAAAGGCTG	360
GATATCAACA	CCAACACCTA	CACATCTCAG	GATCTCAAGA	GTGCACTGGC	AAAATTCAAG	420
GAGGGGGCAG	AGATGGAGAG	TTCAAAGGAA	GACAAGGCAA	GGCAGGCTGA	GGTAAAGCGG	480
CTCTTCCGCC	CCATTGAGGA	ACTGAAGAAA	GACTTTGATG	AGCTGAATGT	TGTCATTGAG	540
ACTGACATGC	AGATCATGGT	ACGGCTGATC	AACAAGTTCA	ATAGTTCCAG	CTCCAGTTTG	600
GAAGAGAAGA	TTGCTGCGCT	CTTTGATCTT	GAATATTATG	TCCATCAGAT	GGACAATGCG	660
CAGGACCTGC	TTTCCTTTGG	TGGTCTTCAA	GTGGTGATCA	ATGGGCTGAA	CAGCACAGAG	720
CCCCTCGTGA	AGGAGTATGC	TGCGTTTGTG	CTGGGCGCTG	CCTTTTCCAG	CAACCCCAAG	780
GTCCAGGTGG	AGGCCATCGA	AGGGGGAGCC	CTGCAGAAGC	TGCTGGTCAT	CCTGGCCACG	840

GAGCAGCCGC	TCACTGCAAA	GAAGAAGGTC	CTGTTTGCAC	TGTGCTCCCT	GCTGCGCCAC	900
TTCCCCATATG	CCCAGCGGCA	GTTCCCTGAAG	CTCGGGGGGC	TGCAGGTCCT	GAGGACCCCTG	960
GTGCAGGAGA	AGGGCACGGA	GGTGCTCGCC	GTGCGCGTGG	TCACACTGCT	CTACGACCTG	1020
GTCACGGAGA	AGATGTTTCGC	CGAGGAGGAG	GCTGAGCTGA	CCCAGGAGAT	GTCCCCAGAG	1080
AAGCTGCAGC	AGTATCGCCA	GGTACACCTC	CTGCCAGGCC	TGTGGGAACA	GGGCTGGTGC	1140
GAGATCACGG	CCCACCTCCT	GGCGCTGCCC	GAGCATGATG	CCCGTGAGAA	GGTGCTGCAG	1200
ACACTGGGCG	TCCTCCTGAC	CACCTGCCGG	GACCGCTACC	GTCAGGACCC	CCAGCTCGGC	1260
AGGACACTGG	CCAGCCTGCA	GGCTGAGTAC	CAGGTGCTGG	CCAGCCTGGA	GCTGCAGGAT	1320
GGTGAGGACG	AGGGCTACTT	CCAGGAGCTG	CTGGGCTCTG	TCAACAGCTT	GCTGAAGGAG	1380
CTGAGATGAG	GCCCCACACC	AGGACTGGAC	TGGGATGCCG	CTAGTGAGGC	TGAGGGGTGC	1440
CAGCGTGGGT	GGGCTTCTCA	GGCAGGAGGA	CATCTTGGA	GTGCTGGCTT	GGCCATTAAA	1500
TGGAAACCTG	AAGGCCAAAA	AAAAAAAAAA	AAAAAAAAAA	AAAAAAAAAA	AAAAAAAAAA	1560
TTCCTGCGGC	CGC					1573

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1185 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GAATTCTGGCA	CGAGGGGGCT	TTAAGGGACA	GCTGAGCCGG	CAGGTGGCAG	ATCAGATGTG	60
GCAGGCTGGG	AAAAGACAAG	CCTCCAGGGC	CTTCAGCTTG	TACGCCAACA	TCGACATCCT	120
CAGACCCTAC	TTTGATGTGG	AGCCTGCTCA	GGTGCGAAGC	AGGCTCCTGG	AGTCCATGAT	180
CCCTATCAAG	ATGGTCAACT	TCCCCAGAA	AATTGCAGGT	GAACCTCTATG	GACCTCTCAT	240
GCTGGTCTTC	ACTCTGGTTG	CTATCCTACT	CCATGGGATG	AAGACGTCTG	ACACTATTAT	300
CCGGGAGGGC	ACCCTGATGG	GCACAGCCAT	TGGCACCTGC	TTCGGCTACT	GGCTGGGAGT	360
CTCATCCTTC	ATTTACTTCC	TTGCCTACCT	GTGCAACGCC	CAGATCACCA	TGCTGCAGAT	420
GTTGGCACTG	CTGGGCTATG	GCCTCTTTGG	GCATTGCATT	GTCCTGTTCA	TCACCTATAA	480
TATCCACCTC	CACGCCCTCT	TCTACCTCTT	CTGGCTGTTG	GTGGGTGGAC	TGTCCACACT	540
GCGCATGGTA	GCAGTGTTGG	TGTCTCGGAC	CGTGGGCCCC	ACACAGCGGC	TGCTCCTCTG	600
TGGCACCCCTG	GCTGCCCTAC	ACATGCTCTT	CCTGCTCTAT	CTGCATTTTG	CCTACCACAA	660

AGTGGTAGAG	GGGATCCTGG	ACACACTGGA	GGGCCCCAAC	ATCCCGCCCA	TCCAGAGGGT	720
CCCCAGAGAC	ATCCCTGCCA	TGCTCCCTGC	TGCTCGGCTT	CCCACCACCG	TCCTCAACGC	780
CACAGCCAAA	GCTGTTGCGG	TGACCCTGCA	GTCACTACTGA	CCCCACCTGA	AATTCTTGGC	840
CAGTCCTCTT	TCCCGCAGCT	GCAGAGAGGA	GGAAGACTAT	TAAAGGACAG	TCCTGATGAC	900
ATGTTTCGTA	GATGGGGTTT	GCAGCTGCCA	CTGAGCTGTA	GCTGCGTAAG	TACCTCCTTG	960
ATGCCTGTCTG	GCACTTCTGA	AAGGCACAAG	GCCAAGAACT	CCTGGCCAGG	ACTGCAAGGC	1020
TCTGCAGCCA	ATGCAGAAAA	TGGGTCAGCT	CCTTTGAGAA	CCCCTCCCCA	CCTACCCCTT	1080
CCTTCCTCTT	TATCTCTCCC	ACATTGTCTT	GCTAAATATA	GACTTGGTAA	TTAAAATGTT	1140
GATTGAAGTC	TGGAATAAAA	AAAAAAAAAA	AATTCCTGCG	GCCGC		1185

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1226 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GAATTCGGCA	CGAGGCAAGC	CACCATCTTC	CTTCGGCCTG	CACCCCTTTA	AAGGCACCCA	60
GACCCCTCTG	GAAAAAGATG	AACTGAAGCC	CTTTGACATC	CTCCAGCCTA	AGGAGTACTT	120
CCAGCTCAGC	CGCCACACGG	TCATTAAGAT	GGGAAGTGAG	AACGAGGCCC	TGGATCTCTC	180
CATGAAGTCA	GTGCCCTGGC	TCAAGGCTGG	TGAAGTCAGT	CCCCCAATCT	TCCAGGAAGA	240
TGCAGCCCTA	GACCTGTCAG	TGGCAGCCCA	CCGGAAATCC	GAGCCTCCCC	CTGAGACACT	300
GTATGACAGT	GGTGCATCAG	TGGACAGCTC	AGGTCACACA	GTGATGGAGA	AACTTCCAG	360
TGGCATGGAA	ATTTCTTTTG	CCCCTGCCAC	GTCCCATGAG	GCCCCAGCCA	TGATGGATAG	420
TCACATCAGC	AGCAGTGATG	CTGCTACCGA	GATGCTCAGC	CAGCCCAACC	ACCCAGCGG	480
CGAAGTCAAG	GCTGAAAATA	ACATTGAGAT	GGTGGGCGAG	TCCCAGGCGG	CCAAGGTCAT	540
TGTCTCTGTC	GAAGATGCTG	TGCCTACCAT	ATTCTGTGGC	AAGATCAAAG	GCCTCTCAGG	600
GGTGTCCACC	AAAAACTTCT	CCTTCAAAAAG	AGAAGACTCC	GTGCTTCAGG	GCTATGACAT	660
CAACAGCCAA	GGGGAAGAGT	CCATGGGAAA	TGCAGAGCCC	CTTAGGAAAC	CCATCAAAAA	720
CCGGAGCATA	AAGTTAAAGA	AAGTGAATC	CCAGGAAGTA	CACATGCTCC	CAATCAAAAA	780
ACAACGGCTG	GCCACCTTTT	TTCCAAGAAA	GTAAATAACG	GCTTTTAAAA	ATTTGTATGA	840
TTATAATATG	GGGAAAGGTG	CATTGGTTTT	ATAAAAAGGC	ATTTAAAAACA	AATTATCTTT	900

GTTAATTATT TTGGGGAGTA GTTGGGAAAT GGAAAGGTGA ATTGGCTCTA GAGGCCCTGT	960
ATGCTAGTAT CATTTTCTTT TTTAATTTTT GACTTTTCAC AAATGAGTAA ATAAGAGCAA	1020
CCTATTTTTC AAGCAGATTG CACATTTTTT GCAGCTTTAA TGGAAATATTG GGTGAATTAG	1080
AGGGGTAAAA AAAGCTATTT TCATTGCCAC AAAGTGCTTT GATGATGTAA TACCTAATAA	1140
AGGGTAGGAT GAATATTTCA CAATAAATGT TTGTTTGCAC TAAAAAAAAA AAAAAAAAAA	1200
AAAAAAAAAA AAATTCCTGC GGCCGC	1226

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1049 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GAATTCGGCA CGAGGGCGCC ATGGTGAAGG TGACGTTCAA CTCCGCTCTG GCCCAGAAGG	60
AGGCCAAGAA GGACGAGCCC AAGAGCGGCG AGGAGGCGCT CATCATCCCC CCCGACGCCG	120
TCGCGGTGGA CTGCAAGGAC CCAGATGATG TGGTACCAGT TGGCCAAAGA AGAGCCTGGT	180
GTTGGTGCAT GTGCTTTGGA CTAGCATTTA TGCTTGCAAG TGTTATTCTA GGAGGAGCAT	240
ACTTGTAACA ATATTTTGCA CTTCAACCAG ATGACGTGTA CTAAGTGGA ATAAAGTACA	300
TCAAAGATGA TGTCATCTTA AATGAGCCCT CTGCAGATGC CCCAGCTGCT CTCTACCAGA	360
CAATTGAAGA AAATATTAAA ATCTTTGAAG AAGAAGAAGT TGAATTTATC AGTGTGCCTG	420
TCCCAGAGTT TGCAGATAGT GATCCTGCCA ACATTGTTCA TGACTTTAAC AAGAACTTA	480
CAGCCTATTT AGATCTTAAC CTGGATAAGT GCTATGTGAT CCCTCTGAAC ACTTCCATTG	540
TTATGCCACC CAGAAACCTA CTGGAGTTAC TTATTAACAT CAAGGCTGGA ACCTATTTGC	600
CTCAGTCCTA TCTGATTCAT GAGCACATGG TTATTACTGA TCGCATTGAA AACATTGATC	660
ACCTGGGTTT CTTTATTTAT CGACTGTGTC ATGACAAGGA AACTTACAAA CTGCAACGCA	720
GAGAAACTAT TAAAGGTATT CAGAAACGTG AAGCCAGCAA TTGTTTCGCA ATTCGGCATT	780
TTGAAAACAA ATTTGCCGTG GAAACTTTAA TTTGTTCTTG AACAGTCAAG AAAACATTA	840
TTGAGGAAAA TTAATATCAC AGCATAACCC CACCCTTTAC ATTTTGTTGC AGTTGATTAT	900
TTTTTAAAGT CTTCTTTTCAT GTAAGTAGCA AACAGGGCTT TACTATCTTT TCATCTCATT	960
AATTCAATTA AAACCATTAC CTTAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA	1020
AAAAAAAAAA AAAAAATTCC TGCGGCCGC	1049

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1142 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GAATTCGGCA CGAGGGGAGA ATACTTTTTG CGATGCCTAC TGGAGACTTT GATTCGAAGC 60  
CCAGTTGGGC CGACCAGGTG GAGGAGGAGG GGGAGGACGA CAAATGTGTC ACCAGCGAGC 120  
TCCTCAAGGG GATCCCTCTG GCCACAGGTG ACACCAGCCC AGAGCCAGAG CTACTGCCGG 180  
GAGCTCCACT GCCGCCTCCC AAGGAGGTCA TCAACGGAAA CATAAAGACA GTGACAGAGT 240  
ACAAGATAGA TGAGGATGGC AAGAAGTTCA AGATTGTCCG CACCTTCAGG ATTGAGACCC 300  
GGAAGGCTTC AAAGGCTGTC GCAAGGAGGA AGAACTGGAA GAAGTTCGGG AACTCAGAGT 360  
TTGACCCCCC CGGACCCAAT GTGGCCACCA CCACTGTCAG TGACGATGTC TCTATGACGT 420  
TCATCACCAG CAAAGAGGAC CTGAACTGCC AGGAGGAGGA GGACCCTATG AACAAATTCA 480  
AGGGCCAGAA GATCGTGTCC TGCCGCATCT GCAAGGGCGA CCACTGGACC ACCCGCTGCC 540  
CCTACAAGGA TACGCTGGGG CCCATGCAGA AGGAGCTGGC CGAGCAGCTG GGCCTGTCTA 600  
CTGGCGAGAA GGAGAAGCTG CCGGGAGAGC TAGAGCCGGT GCAGGCCACG CAGAACAAGA 660  
CAGGGAAGTA TGTGCCGCCG AGCCTGCGCG ACGGGGCCAG CCGCCGCGGG GAGTCCATGC 720  
AGCCCAACCG CAGAGCCGAC GACAACGCCA CCATCCGTGT CACCAACTTG CGCAGAGGAC 780  
ACGCGTGAGA CCGACCTGCA GGAGCTCTTC CGGCCTTTTG GCTCCATCTC CCGCATCTAC 840  
CTGGCTAAGG ACAAGACCAC TGGCCAATCC AAGGGCTTTG CCTTCATCAG CTTCACCCGC 900  
CGCGAGGATG CTGCGCGTGC CATTGCCGGG GTGTCCGGCT TTGGCTACGA CCACCTCATC 960  
CTCAACGTCG AGTGGGCCAA GCCGTCCACC AACTAAGCCA GCTGCCACTG TGTACTCGGT 1020  
CCGGGACCCCT TGGCGACAGA AGACAGCCTC CGAGAGCGCG GGCTCCAAGG GCAATAAAGC 1080  
AGCTCCACTC TCAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAAAT TCCTGCGGCC 1140  
GC 1142

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1696 base pairs

(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GAATTCGGCA	CGAGGGAAAC	ATGGCGGTAG	GCTGGGACCA	TAACACAAGC	ATGACTATAT	60
GAAGGAAGAG	GAAGGTTTTCT	CTGAAGATGA	GGCGACTGAA	TCGGAAAAAA	ACTTTAAGTT	120
TGGTAAAAGA	GTTGGATGCC	TTTCCGAAGG	TTCTGAGAG	CTATGTAGAG	ACTTCAGCCA	180
GTGGAGGTAC	AGTTTCTCTA	ATAGCATTTA	CAACTATGGC	TTTATTAACC	ATAATGGAAT	240
TCTCAGTATA	TCAAGATACA	TGGATGAAGT	ATGAATACGA	AGTAGACAAG	GATTTTTCTA	300
GCAAATTAAG	AATTAATATA	GATATTACTG	TTGCCATGAA	GTGTCAATAT	GTTGGAGCGG	360
ATGTATTGGA	TTTAGCAGAA	ACAATGGTTG	CATCTGCAGA	TGGTTTAGTT	TATGAACCAA	420
CAGTATTTGA	TCTTTCACCA	CAGCAGAAAG	AGTGGCAGAG	GATGCTGCAG	CTGATTCAGA	480
GTAGGCTACA	AGAAGAGCAT	TCACCTCAAG	ATGTGATATT	TAAAAGTGCT	TTTAAAAGTA	540
CATCAACAGC	TCTTCCACCA	AGAGAAGATG	ATTCATCACA	GTCTCCAAAT	GCATGCAGAA	600
TTCATGGCCA	TCTATATGTC	AATAAAGTAG	CAGGGAATTT	TCACATAACA	GTGGGCAAGG	660
CAATTCACCA	TCCTCGTGGT	CATGCACATT	TGGCAGCACT	TGTCAACCAT	GAATCTTACA	720
ATTTTCTCA	TAGAATAGAT	CATTTGTCTT	TTGGAGAGCT	TGTTCCAGCA	ATTATTAATC	780
CTTTAGATGG	AACTGAAAAA	ATTGCTATAG	ATCACAACCA	GATGTTCCAA	TATTTTATTA	840
CAGTTGTGCC	AACAAAACTA	CATACATATA	AAATATCAGC	AGACACCCAT	CAGTTTTCTG	900
TGACAGAAAG	GGAACGTATC	ATTAACCATG	CTGCAGGCAG	CCATGGAGTC	TCTGGGATAT	960
TTATGAAATA	TGATCTCAGT	TCTCTTATGG	TGACAGTTAC	TGAGGAGCAC	ATGCCATTCT	1020
GGCAGTTTTT	TGTAAGACTC	TGTGGTATTG	TTGGAGGAAT	CTTTTCAACA	ACAGGCATGT	1080
TACATGGAAT	TGGAAAATTT	ATAGTTGAAA	TAATTTGCTG	TCGTTTCAGA	CTTGGATCCT	1140
ATAAACCTGT	CAATTCTGTT	CCTTTTGAGG	ATGGCCACAC	AGACAACCAC	TTACCTCTTT	1200
TAGAAAATAA	TACACATTAA	CACCTCCCGA	TTGAAGGAGA	AAAACTTTTT	GCCTGAGACA	1260
TAAAACCTTT	TTTTAATAAT	AAAATATTGT	GCAATATATT	CAAAGAAAAG	AAAACACAAA	1320
TAAGCAGAAA	ACATACTTAT	TTAAAAAAG	AAAAAAAAGG	ATAAAAAAAC	CCAAACTGAA	1380
ATTCTATATA	CGTTGTGTCT	GTTACAAATG	TCGTAGAAGA	AATCATGCAG	CTAAACGATG	1440
AAGAAGCCCA	ACTGGAGTGT	TGCTTTGAAG	ATGACGCCTT	CTTATATTTT	CATAGCAAAT	1500
GGGTGGTATC	AAAATCAGAC	ATTGCTTCTT	GCTGATAAAA	AGCCTGAAGG	AAATAAGTGA	1560
AACTACATCT	ATGGGAAAAA	AAAAAACATT	GAGAAGTGCA	AATGTTGCA	TCCTTTTGTT	1620
TTTAAAAGAT	ATGATGTCAG	AATAAAATGT	GGAAAACATA	CGGAAAAAAA	AAAAAAAATA	1680
AAATTCCTGC	GGCCGC					1696

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1100 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GAATTCGGCA	CGAGGCGGCA	CGAGGCGGCA	CGAGGGTGGC	ATATCACGGC	CATGGGGTCT	60
CAGCATTCCG	CTGCTGCTCG	CCCCTCCTCC	TGCAGGCGAA	AGCAAGAAGA	TGACAGGGAC	120
GGTTTGCTGG	CTGAACGAGA	GCAGGAAGAA	GCCATTGCTC	AGTTCCTATA	TGTGGAATTC	180
ACCGGGAGAG	ATAGCATCAC	CTGTCTCACG	TGCCAGGGGA	CAGGCTACAT	TCCAACAGAG	240
CAAGTAAATG	AGTTGGTGGC	TTTGATCCCA	CACAGTGATC	AGAGATTGCG	CCCTCAGCGA	300
ACTAAGCAAT	ATGTCCTCCT	GTCCATCCTG	CTTTGTCTCC	TGGCATCTGG	TTTGGTGGTT	360
TTCTTCCTGT	TTCCGCATTC	AGTCCTTGTG	GATGATGACG	GCATCAAAGT	GGTGAAAGTC	420
ACATTTAATA	AGCAAGACTC	CCTTGTAATT	CTCACCATCA	TGGCCACCCT	GAAAATCAGG	480
AACTCCAACT	TCTACACGGT	GGCAGTGACC	AGCCTGTCCA	GCCAGATTCA	GTACATGAAC	540
ACAGTGGTCA	GTACATATGT	GACTACTAAC	GTCTCCCTTA	TTCCACCTCG	GAGTGAGCAA	600
CTGGTGAATT	TTACCGGGAA	GGCCGAGATG	GGAGGACCGT	TTTCCTATGT	GTACTTCTTC	660
TGCACGGTAC	CTGAGATCCT	GGTGCACAAC	ATAGTGATCT	TCATGCGAAC	TTCAGTGAAG	720
ATTTCATACA	TTGGCCTCAT	GACCCAGAGC	TCCTTGAGAG	CACATCACTA	TGTGGATTGT	780
GGAGGAAATT	CCACAGCTAT	TTAACAACCTG	CTATTGGTTC	TTCCACACAG	CGCCTGTAGA	840
AGAGAGCACA	GCATATGTTC	CCAAGGCCTG	AGTTCTGGAC	CTACCCCCAC	GTGGTGTAAAG	900
CAGAGGAGGA	ATTGGTTCAC	TTAACTCCCA	GCAAACATCC	TCCTGCCACT	TAGGAGGAAA	960
CACCTCCCTA	TGGTACCATT	TATGTTTCTC	AGAACCAGCA	GAATCAGTGC	CTAGCCTGTG	1020
CCCAGCAAAT	AGTTGGCACT	CAATAAAGAT	TTGCAGAATT	TAAAAAAAAA	AAAAAAAAAA	1080
AAAAAAATTC	CTGCGGCCGC					1100

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1588 base pairs
- (B) TYPE: nucleic acid



(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GAATTCGGCA	CGAGGGTACC	TGCTTTTCTA	TTGCCTCTTT	GAAACAATGG	TCACGTGTTT	60
CCATGTTCCC	TACTCGGCTC	TCACCATGTT	CATCAGCACC	GAGCAGACTG	AGCGGGATTC	120
TGCCACCGCC	TATCGGATGA	CTGTGGAAGT	GCTGGGCACA	GTGCTGGGCA	CGGCGATCCA	180
GGGACAAATC	GTGGGCCAAG	CAGACACGCC	TTGTTTCCAG	GACCTCAATA	GCTCTACAGT	240
AGCTTCACAA	AGTGCCAACC	ATACACATGG	CACCACCTCA	CACAGGGAAA	CGCAAAAGGC	300
ATACCTGCTG	GCAGCGGGGG	TCATTGTCTG	TATCTATATA	ATCTGTGCTG	TCATCCTGAT	360
CCTGGGCGTG	CGGGAGCAGA	GAGAACCCTA	TGAAGCCCAG	CAGTCTGAGC	CAATCGCCTA	420
CTTCCGGGGC	CTACGGCTGG	TCATGAGCCA	CGGCCCATAC	ATCAAACCTA	TTACTGGCTT	480
CCTCTTCACC	TCCTTGGCTT	TCATGCTGGT	GGAGGGGAAC	TTTGTCTTGT	TTTGCACCTA	540
CACCTTGGGC	TTCCGCAATG	AATTCCAGAA	TCTACTCCTG	GCCATCATGC	TCTCGGCCAC	600
TTTAACCATT	CCCATCTGGC	AGTGGTTCTT	GACCCGGTTT	GGCAAGAAGA	CAGCTGTATA	660
TGTTGGGATC	TCATCAGCAG	TGCCATTTCT	CATCTTGGTG	GCCCTCATGG	AGAGTAACCT	720
CATCATTACA	TATGCGGTAG	CTGTGGCAGC	TGGCATCAGT	GTGGCAGCTG	CCTTCTTACT	780
ACCCTGGTCC	ATGCTGCCTG	ATGTCATTGA	CGACTTCCAT	CTGAAGCAGC	CCCACTTCCA	840
TGGAACCGAG	CCCATCTTCT	TCTCCTTCTA	TGTCTTCTTC	ACCAAGTTTG	CCTCTGGAGT	900
GTCACTGGGC	ATTTCTACCC	TCAGTCTGGA	CTTTGCAGGG	TACCAGACCC	GTGGCTGCTC	960
GCAGCCGGAA	CGTGTCAAGT	TTACACTGAA	CATGCTCGTG	ACCATGGCTC	CCATAGTTCT	1020
CATCCTGCTG	GGCCTGCTGC	TCTTCAAAAT	GTACCCCAT	GATGAGGAGA	GGCGGCGGCA	1080
GAATAAGAAG	GCCCTGCAGG	CACTGAGGGA	CGAGGCCAGC	AGCTCTGGCT	GCTCAGAAAC	1140
AGACTCCACA	GAGCTGGCTA	GCATCCTCTA	GGGCCCCCCA	CGTTGCCCCG	AGCCACCATG	1200
CAGAAGGCCA	CAGAAGGGAT	CAGGACCTGT	CTGCCGGCTT	GCTGAGCAGC	TGGACTGCAG	1260
GTGCTAGGAA	GGGAAGTGAA	GACTCAAGGA	GGTGGCCCAG	GACACTTGCT	GTGCTCACTG	1320
TGGGGGCGGC	TGCTCTGTGG	CCTCCTGCCT	CCCCTCTGCC	TGCCTGTGGG	GCCAAGCCCT	1380
GGGGCTGCCA	CTGTGAATAT	GCCAAGGACT	GATCGGGCCT	AGCCCGGAAC	ACTAATGTAG	1440
AAACCTTTTT	TTTACAGAGC	CTAATTAATA	ACTTAATGAC	TGTGTACATA	GCAATGTGTG	1500
TGTATGTATA	TGTCTGTGAG	CTATTAATGT	TATTAATTTT	CATAAAAGCT	GGAAAGCAAA	1560
AAAAAAAAAA	AAAAATTCCT	GCGGCCGC				1588

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1535 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GAATTCGGCA	CGAGGCGGAA	GTCCCGTCTC	ACGTTTGCCC	TGGCAGCGCG	CGAGGCTGGT	60
GAGTCGGCAG	CCCTGTGGCA	GCCGGCGGGC	TGGTTTCCAT	GGTTGCACGA	TTAGGAACCA	120
CCAGCTGCTG	CATCCCATGG	CCAGGGGTGG	CGTCCAGGTG	GCAGAGCAGC	TAGGAACGCA	180
AGGCCTGAAC	CTGGGGCCAG	ACACCCTGCT	CTCCCGGCCA	TGGTCAACGA	CCCTCCAGTA	240
CCTGCCTTAC	TGTGGGCCCA	GGAGGTGGGC	CAAGTCTTGG	CAGGCCGTGC	CCGCAGGCTG	300
CTGCTGCAGT	TTGGGGTGCT	CTTCTGCACC	ATCCTCCTTT	TGCTCTGGGT	GTCTGTCTTC	360
CTCTATGGCT	CCTTCTACTA	TTCCTATATG	CCGACAGTCA	GCCACCTCAG	CCCTGTGCAT	420
TTCTACTACA	GGACCGACTG	TGATTCTCTC	ACCACCTCAC	TCTGCTCCTT	CCCTGTTGCC	480
AATGTCTCGC	TGACTAAGGG	TGGACGTGAT	CGGGTGCTGA	TGTATGGACA	GCCGTATCGT	540
GTTACCTTAG	AGCTTGAGCT	GCCAGAGTCC	CCTGTGAATC	AAGATTTGGG	CATGTTCTTG	600
GTCACCATTT	CCTGCTACAC	CAGAGGTGGC	CGAATCATCT	CCACTTCTTC	GCGTTCGGTG	660
ATGCTGCATT	ACCGCTCAGA	CCTGCTCCAG	ATGCTGGACA	CACTGGTCTT	CTCTAGCCTC	720
CTGCTATTTG	GCTTTGCAGA	GCAGAAGCAG	CTGCTGGAGG	TGGAACCTTA	CGCAGACTAT	780
AGAGAGAACT	CGTACGTGCC	GACCACTGGA	GCGATCATTG	AGATCCACAG	CAAGCGCATC	840
CAGCTGTATG	GAGCCTACCT	CCGCATCCAC	GCGCACTTCA	CTGGGCTCAG	ATACCTGCTA	900
TACAACTTCC	CGATGACCTG	CGCCTTCATA	GGTGTGTTG	GCAACTTCAC	CTTCCTCAGC	960
GTCATCGTGC	TCTTCAGCTA	CATGCAGTGG	GTGTGGGGGG	GCATCTGGCC	CCGACACCGC	1020
TTCTCTTTGC	AGGTTAACAT	CCGAAAAAGA	GACAATTCCC	GGAAGGAAGT	CCAACGAAGG	1080
ATCTCTGCTC	ATCAGCCAGG	GCCTGAAGGC	CAGGAGGAGT	CAACTCCGCA	ATCAGATGTT	1140
ACAGAGGATG	GTGAGAGCCC	TGAAGATCCC	TCAGGGACAG	AGGTCAGCTG	TCCGAGGAGG	1200
AGAAACCAGA	TCAGCAGCCC	CTGAGCGGAG	AAGAGGAGCT	AGAGCCTGAG	GCCAGTGATG	1260
GTTCAGGCTC	CTGGGAAGAT	GCAGCTTTGC	TGACGGAGGC	CAACCTGCCT	GCTCCTGCTC	1320
CTGCTTCTGC	TTCTGCCCC	GTCTTAGAGA	CTCTGGGCAG	CTCTGAACCT	GCTGGGGGTG	1380
CTCTCCGACA	GCGCCCCACC	TGCTCTAGTT	CCTGAAGAAA	AGGGGCAGAC	TCCTCACATT	1440
CCAGCACTTT	CCCACCTGAC	TCCTCTCCCC	TCGTTTTTCC	TTCAATAAAC	TATTTTGTGT	1500
CAAAAAAAAA	AAAAAAAAAA	AATTCCTGCG	GCCGC			1535

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1322 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GAATTCGGCA	CGAGGGCGGG	CGCTACGGGC	TTGACTCCCC	CAAGGCCGAG	GTCCGCGGCC	60
AGGTGCTGGC	GCCGCTGCCC	CTCCACGGAG	TTGCTGATCA	TCTGGGCTGT	GATCCACAAA	120
CCCGGTTCTT	TGTCCCTCCT	AATATCAAAC	AGTGGATTGC	CTTGCTGCAG	AGGGGAAACT	180
GCACGTTTAA	AGAGAAAATA	TCACGGGCCG	CTTCCACAA	TGCAGTTGCT	GTAGTCATCT	240
ACAATAATAA	ATCCAAAGAG	GAGCCAGTTA	CCATGACTCA	TCCAGGCACT	GGAGATATTA	300
TTGCTGTCAT	GATAACAGAA	TTGAGGGGTA	AGGATATTTT	GAGTTATCTG	GAGAAAAACA	360
TCTCTGTACA	AATGACAATA	GCTGTTGGAA	CTCGAATGCC	ACCGAAGAAC	TTCAGCCGTG	420
GCTCTCTAGT	CTTCGTGTCA	ATATCCTTTA	TTGTTTTGAT	GATTATTTCT	TCAGCATGGC	480
TCATATTCTA	CTTCATTCAA	AAGATCAGGT	ACACAAATGC	ACGCGACAGG	AACCAGCGTC	540
GTCTCGGAGA	TGCAGCCAAG	AAAGCCATCA	GTAAATTGAC	AACCAGGACA	GTAAAGAAGG	600
GTGACAAGGA	AACTGACCCA	GACTTTGATC	ATTGTGCAGT	CTGCATAGAG	AGCTATAAGC	660
AGAATGATGT	CGTCCGAATT	CTCCCCTGCA	AGCATGTTTT	CCACAAATCC	TGCGTGATC	720
CCTGGCTTAG	TGAACATTGT	ACCTGTCCTA	TGTGCAAACT	TAATATATTG	AAGGCCCTGG	780
GAATTGTGCC	GAATTTGCCA	TGTACTGATA	ACGTAGCATT	CGATATGGAA	AGGCTCACCA	840
GAACCCAAGC	TGTTAACCGA	AGATCAGCCC	TCGGCGACCT	CGCCGGCGAC	AACTCCCTTG	900
GCCTTGAGCC	ACTTCGAACT	TCGGGGATCT	CACCTCTTCC	TCAGGATGGG	GAGCTCACTC	960
CGAGAACAGG	AGAAATCAAC	ATTGCAGTAA	CAAAAGAATG	GTTTATTATT	GCCAGTTTTG	1020
GCCTCCTCAG	TGCCCTCACA	CTCTGCTACA	TGATCATCAG	AGCCACAGCT	AGCTTGAATG	1080
CTAATGAGGT	AGAATGGTTT	TGAAGAAGAA	AAAACCTGCT	TTCTGACTGA	TTTTGCCTTG	1140
AAGGAAAAAA	GAACCTATTT	TTGTGCATCA	TTTACCAATC	ATGCCACACA	AGCATTTATT	1200
TTTAGTACAT	TTTATTTTTT	CATAAAATTG	CTAATGCCAA	AGCTTTGTAT	TAAAAGAAAT	1260
AAATAATAAA	ATAAAAAAAA	AAAAAATAAA	AAAAAATAAA	AAAAAATAAT	TCCTGCGGCC	1320
GC						1322

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1711 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GAATTCGGCA	CGAGGCCCTC	CCGCGCTCCC	GGGGCGCGCG	GGCCGCGCCC	CCGACGCCCT	60
ACATATACTC	AGGTGCGCCC	CACCTGTCCG	CCCGCACCTG	CTGGCTCACC	TCCGAGCCAC	120
CTCTGCTGCG	CACCGCAGCC	TCGGACCTAC	AGCCCAGGAT	ACTTTGGGAC	TTGCCGGCGC	180
TCAGAAACGC	GCCCAGACGG	CCCCTCCACC	TTTTGTTTGC	CTAGGGTCGC	CGAGAGCGCC	240
CGGAGGGAAC	CGCCTGGCCT	TCGGGGACCA	CCAATTTTGT	CTGGAACCAC	CCTCCCGGCG	300
TATCCTACTC	CCTGTGCCGC	GAGGCCATCG	CTTCACTGGA	GGGGTCGATT	TGTGTGTAGT	360
TTGGTGACAA	GATTTGCATT	CACCTGGCCC	AAACCCTTTT	TGTCTCTTTG	GGTGACCGGA	420
AAACTCCACC	TCAAGTTTTC	TTTTGTGGGG	CTGCCCCCA	AGTGTGTTTT	GTTTTACTGT	480
AGGGTCTCCC	GCCCCGCGCC	CCCAGTGTTT	TCTGAGGGCG	GAAATGGCCA	ATTCGGGCCT	540
GCAGTTGCTG	GGCTTCTCCA	TGGCCCTGCT	GGGCTGGGTG	GGTCTGGTGG	CCTGCACCGC	600
CATCCCGCAG	TGGCAGATGA	GCTCCTATGC	GGGTGACAAC	ATCATCACGG	CCCAGGCCAT	660
GTACAAGGGG	CTGTGGATGG	ACTGCGTCAC	GCAGAGCACG	GGGATGATGA	GCTGCAAAAT	720
GTACGACTCG	GTGCTCGCCC	TGTCCGCGGC	CTTGACGGCC	ACTCGAGCCC	TAATGGTGGT	780
CTCCCTGGTG	CTGGGCTTCC	TGGCCATGTT	TGTGGCCACG	ATGGGCATGA	AGTGCACGCG	840
CTGTGGGGGA	GACGACAAAG	TGAAGAAGGC	CCGTATAGCC	ATGGGTGGAG	GCATAATTTT	900
CATCGTGCCA	GGTCTTGCCG	CCTTGGTAGC	TTGCTCCTGG	TATGGCCATC	AGATTGTCAC	960
AGACTTTTAT	AACCCTTTGA	TCCCTACCAA	CAATTAAGTAT	GAGTTTGGCC	CTGCCATCTT	1020
TATTGGCTGG	GCAGGGTCTG	CCCTAGTCAT	CCTGGGAGGT	GCACTGCTCT	CCTGTTCCCTG	1080
TCCTGGGAAT	GAGAGCAAGG	CTGGGTACCG	TGCACCCCGC	TCTTACCCTA	AGTCCAACTC	1140
TTCCAAGGAG	TATGTGTGAC	CTGGGATCTC	CTTGCCCCAG	CCTGACAGGC	TATGGGAGTG	1200
TCTAGATGCC	TGAAAGGGCC	TGGGGCTGAG	CTCAGCCTGT	GGGCAGGGTG	CCGGACAAAG	1260
GCCTCCTGGT	CACTCTGTCC	CTGCACTCCA	TGTATAGTCC	TCTTGGGTTG	GGGGTGGGGG	1320
GGTGCCGTTG	GTGGGAGAGA	CAAAAAGAGG	GAGAGTGTGC	TTTTTGTACA	GTAATAAAAA	1380
ATAAGTATTG	GGAAGCAGGC	TTTTTTCCCT	TCAGGGCCTC	TGCTTTCCTC	CCGTCCAGAT	1440
CCTTGACAGG	AGCTTGAAC	CTTAGTGCAC	CTACTTCAGT	TCAGAACACT	TAGCACCCCA	1500
CTGACTCCAC	TGACAATTGA	CTAAAAGATG	CAGGTGCTCG	TATCTCGACA	TTCATTCCCA	1560
CCCCCTCTT	ATTAAATAG	CTACCAAAGT	ACTTCTTTTT	TAATAAAAAA	ATAAAGATTT	1620

TTATTAGGTA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 1680  
 AAAAAAAAAA AAAAAAATT CCTGCGGCCG C 1711

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1553 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GAATTCGGCA	CGAGGGCAGG	TCCAGAGTAA	AGTCACTGAA	GAGTGGAAGC	GAGGAAGGAA	60
CAGGATGATT	AGACCTCAGC	TGCGGACCGC	GGGGCTGGGA	CGATGCCTCC	TGCCGGGGCT	120
GCTGCTGCTC	CTGGTGCCCG	TCCTCTGGGC	CGGGGCTGAA	AAGCTACATA	CCCAGCCCTC	180
CTGCCCCGCG	GTCTGCCAGC	CCACGCGCTG	CCCCGCGCTG	CCCACCTGCG	CGCTGGGGAC	240
CACGCCGGTG	TTCGACCTGT	GCCGCTGTTG	CCGCGTCTGC	CCCGCGGCCG	AGCGTGAAGT	300
CTGCGGCGGG	GCGCAGGGCC	AACCGTGCGC	CCCGGGGCTG	CAGTGCCTCC	AGCCGCTGCG	360
CCCCGGGTTC	CCCAGCACCT	GCGGTTGCCC	GACGCTGGGA	GGGGCCGTGT	GCGGCAGCGA	420
CAGGCGCACC	TACCCACAGC	TGTGCGCGCT	CCGGGCCGAA	AACCGCGCCG	CGCGCCGCCT	480
GGGCAAGGTC	CCGGCCGTGC	CTGTGCAGTG	GGGGAAGTGC	GGGGATACAG	GGACCAGAAG	540
CGCAGGCCCC	CTCAGGAGGA	ATTACAACCT	CATCGCCGCG	GTGGTGGAGA	AGGTGGCGCC	600
ATCGGTGGTT	CACGTGCAGC	TGTGGGGCAG	GTTACTTCAC	GGCAGCAGGC	TTGTTCCTGT	660
GTACAGTGGC	TCTGGGTTCA	TAGTGTCTGA	GGACGGGCTC	ATTATTACCA	ATGCCCATGT	720
TGTCAGGAAC	CAGCAGTGGA	TTGAGGTGGT	GCTCCAGAAT	GGGGCCCGTT	ATGAAGCTGT	780
TGTCAAGGAT	ATTGACCTTA	AATTGGATCT	TGCGGTGATT	AAGATTGAAT	CAAATGCTGA	840
ACTTCCTGTA	CTGATGCTGG	GAAGATCATC	TGACCTTCGG	GCTGGAGAGT	TTGTGGTGGC	900
TTTGGGCAGC	CCATTTTCTC	TGCAGAACAC	AGCTACTGCA	GGAATTGTCA	GCACCAAACA	960
GCGAGGGGGC	AAAGAACTGG	GGATGAAGGA	TTCAGATATG	GACTACGTCC	AGATTGATGC	1020
CACAATTAAC	TATGGGAATT	CTGGTGGTCC	TCTGGTGAAC	TTGGATGGTG	ATGTGATTGG	1080
CGTCAATTCA	TTGAGGGTGA	CTGATGGAAT	CTCCTTTGCA	ATTCCTTCAG	ATCGAGTTAG	1140
GCAGTTCTTG	GCAGAAATAC	ATGAGCACCA	GATGAAAGGA	AAGGCGTTTT	CAAATAAGAA	1200
ATATCTGGGT	CTGCAAATGC	TGTCCCTCAC	TGTGCCCCCT	AGTGAAGAAT	TGAAAATGCA	1260
TTATCCAGAT	TTCCCTGATG	TGAGTTCTGG	GGTTTATGTA	TGTAAAGTGG	TTGAAGGAAC	1320

AGCTGCTCAA AGCTCTGGAT TGAGAGATCA CGATGTAATT GTCAACATAA ATGGGAAACC	1380
TATTACTACT ACAACTGATG TTGTTAAAGC TCTTGACAGT GATTCCCTTT CCATGGCTGT	1440
TCTTCGGGGA AAAGATAATT TGCTCCTGAC AGTCATACCT GAAACAATCA ATTAAATATC	1500
TTGTTTTTAAA GTGGGATTAT CTAAAAAAA AAAAAAAAAA TTCCTGCGGC CGC	1553

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1596 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GAATTCGGCA CGAGGGGAGC CGCTCCCGGA GCCCGGCCGT AGAGGCTGCA ATCGCAGCCG	60
GGAGCCCGCA GCCCGCGCCC CGAGCCCGCC GCCGCCCTTC GAGGGCGCCC CAGGCCGCGC	120
CATGGTGAAG GTGACGTTCA ACTCCGCTCT GGCCCAAG GAGGCCAAGA AGGACGAGCC	180
CGAGAGCGGC GAGGAGGCGC TCATCATCCC CCCCAGCGCC GTCGCGGTGG ACTGCAAGGA	240
CCCAGATGAT GTGGTACCAG TTGGCCAAAG AAGAGCCTGG TGTGGTGCA TGTGCTTTGG	300
ACTAGCATTT ATGCTTGCA GTGTTATTCT AGGAGGAGCA TACTTGTAACA AATATTTTGC	360
ACTTCAACCA GATGACGTGT ACTACTGTGG AATAAAGTAC ATCAAAGATG ATGTCATCTT	420
AAATGAGCCC TCTGCAGATG CCCCAGCTGC TCTCTACCAG ACAATTGAAG AAAATATTAA	480
AATCTTTGAA GAAGAAGAAG TTGAATTTAT CAGTGTGCCT GTCCCAGAGT TTGCAGATAG	540
TGATCCTGCC AACATTGTTC ATGACTTTAA CAAGAACTT ACAGCCTATT TAGATCTTAA	600
CCTGGATAAG TGCTATGTGA TCCCTCTGAA CACTTCCATT GTTATGCCAC CCAGAAACCT	660
ACTGGAGTTA CTTATTACA TCAAGGCTGG AACCTATTTG CCTCAGTCCT ATCTGATTCA	720
TGAGCACATG GTTATTACTG ATCGCATTGA AAACATTGAT CACCTGGGTT TCTTTATTTA	780
TCGACTGTGT CATGACAAGG AACTTACAA ACTGCAACGC AGAGAAACTA TTAAAGGTAT	840
TCAGAAACGT GAAGCCAGCA ATTGTTTCGC AATTCGGCAT TTTGAAAACA AATTTGCCGT	900
GGAACTTTA ATTTGTTCTT GAACAGTCAA GAAAAACATT ATTGAGGAAA ATTAATATCA	960
CAGCATAACC CCACCTTTA CATTTTGTGC AGTGATATTT TTTAAAGTCT CTTTCATGTA	1020
AGTAGCAAAC AGGGCTTTAC TATCTTTTCA TCTCATTAAT TCAATTAAAA CCATTACCTT	1080
AAAATTTTTT TCTTTCGAAG TGTGGTGTCT TTTATATTTG AATTAGTAAC TGTATGAAGT	1140

CATAGATAAT AGTACATGTC ACCTTAGGTA GTAGGAAGAA TTACAATTTC TTAAATCAT	1200
TTATCTGGAT TTTTATGTTT TATTAGCATT TTCAAGAAGA CGGATTATCT AGAGAATAAT	1260
CATATATATG CATACGTAAA AATGGACCAC AGTGACTTAT TTGTAGTTGT TAGTTGCCCT	1320
GCTACCTAGT TTGTTAGTGC ATTTGAGCAC ACATTTTAAAT TTCCTCTAA TTAAAATGTG	1380
CAGTATTTTC AGTGTCAAAT ATATTTAACT ATTTAGAGAA TGATTTCAC CTTTATGTTT	1440
TAATATCCTA GGCATCTGCT GTAATAATAT TTTAGAAAAT GTTTGGAATT TAAGAAATAA	1500
CTTGTGTTAC TAATTTGTAT AACCCATATC TGTGCAATGG AATATAAATA TCACAAAGTT	1560
GTTTAAAAAA AAAAAAAAAA AAATTCCTGC GGCCGC	1596

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 400 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Met	Ala	Trp	Arg	Arg	Arg	Glu	Ala	Gly	Val	Gly	Ala	Arg	Gly	Val	Leu
1				5					10					15	
Ala	Leu	Ala	Leu	Leu	Ala	Leu	Ala	Leu	Cys	Val	Pro	Gly	Ala	Arg	Gly
			20						25					30	
Arg	Ala	Leu	Glu	Trp	Phe	Ser	Ala	Val	Val	Asn	Ile	Glu	Tyr	Val	Asp
			35						40					45	
Pro	Gln	Thr	Asn	Leu	Thr	Val	Trp	Ser	Val	Ser	Glu	Ser	Gly	Arg	Phe
			50						55					60	
Gly	Asp	Ser	Ser	Pro	Lys	Glu	Gly	Ala	His	Gly	Leu	Val	Gly	Val	Pro
65					70					75				80	
Trp	Ala	Pro	Gly	Gly	Asp	Leu	Glu	Gly	Cys	Ala	Pro	Asp	Thr	Arg	Phe
			85						90					95	
Phe	Val	Pro	Glu	Pro	Gly	Gly	Arg	Gly	Ala	Ala	Pro	Trp	Val	Ala	Leu
			100						105					110	
Val	Ala	Arg	Gly	Gly	Cys	Thr	Phe	Lys	Asp	Lys	Val	Leu	Val	Ala	Ala

115	120	125
Arg Arg Asn Ala Ser Ala Val Val Leu Tyr Asn Glu Glu Arg Tyr Gly		
130	135	140
Asn Ile Thr Leu Pro Met Ser His Ala Gly Thr Gly Asn Ile Val Val		
145	150	155
Ile Met Ile Ser Tyr Pro Lys Gly Arg Glu Ile Leu Glu Leu Val Gln		
165	170	175
Lys Gly Ile Pro Val Thr Met Thr Ile Gly Val Gly Thr Arg His Val		
180	185	190
Gln Glu Phe Ile Ser Gly Gln Ser Val Val Phe Val Ala Ile Ala Phe		
195	200	205
Ile Thr Met Met Ile Ile Ser Leu Ala Trp Leu Ile Phe Tyr Tyr Ile		
210	215	220
Gln Arg Phe Leu Tyr Thr Gly Ser Gln Ile Gly Ser Gln Ser His Arg		
225	230	235
Lys Glu Thr Lys Lys Val Ile Gly Gln Leu Leu Leu His Thr Val Lys		
245	250	255
His Gly Glu Lys Gly Ile Asp Val Asp Ala Glu Asn Cys Ala Val Cys		
260	265	270
Ile Glu Asn Phe Lys Val Lys Asp Ile Ile Arg Ile Leu Pro Cys Lys		
275	280	285
His Ile Phe His Arg Ile Cys Ile Asp Pro Trp Leu Leu Asp His Arg		
290	295	300
Thr Cys Pro Met Cys Lys Leu Asp Val Ile Lys Ala Leu Gly Tyr Trp		
305	310	315
Gly Glu Pro Gly Asp Val Gln Glu Met Pro Ala Pro Glu Ser Pro Pro		
325	330	335
Gly Arg Asp Pro Ala Ala Asn Leu Ser Leu Ala Leu Pro Asp Asp Asp		
340	345	350
Gly Ser Asp Asp Ser Ser Pro Pro Ser Ala Ser Pro Ala Glu Ser Glu		
355	360	365
Pro Gln Cys Asp Pro Ser Phe Lys Gly Asp Ala Gly Glu Asn Thr Ala		
370	375	380
Leu Leu Glu Ala Gly Arg Ser Asp Ser Arg His Gly Gly Pro Ile Ser		
385	390	395
		400



(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 291 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Met Asp Lys Gly Ser Ala Gly His Pro Gly Gly Val Leu Val Trp Gly  
1 5 10 15  
Arg Ser Pro Ala Pro Thr Ala Leu Trp Gly Ala Ser Pro Trp Leu Ser  
20 25 30  
Pro Leu Thr Ser Ala Leu Arg Gln Pro Leu His Arg Ala Pro Leu Leu  
35 40 45  
Pro Gly Gln Leu Cys Trp Ser Pro Arg Pro Leu Glu Lys Asn Lys Ala  
50 55 60  
Met Gly Arg Pro Leu Leu Leu Pro Leu Leu Leu Leu Leu Gln Pro Pro  
65 70 75 80  
Ala Phe Leu Gln Pro Gly Gly Ser Thr Gly Ser Gly Pro Ser Tyr Leu  
85 90 95  
Tyr Gly Val Thr Gln Pro Lys His Leu Ser Ala Ser Met Gly Gly Ser  
100 105 110  
Val Glu Ile Pro Phe Ser Phe Tyr Tyr Pro Trp Glu Leu Ala Ile Val  
115 120 125  
Pro Asn Val Arg Ile Ser Trp Arg Arg Gly His Phe His Gly Gln Ser  
130 135 140  
Phe Tyr Ser Thr Arg Pro Pro Ser Ile His Lys Asp Tyr Val Asn Arg  
145 150 155 160  
Leu Phe Leu Asn Trp Thr Glu Gly Gln Glu Ser Gly Phe Leu Arg Ile  
165 170 175  
Ser Asn Leu Arg Lys Glu Asp Gln Ser Val Tyr Phe Cys Arg Val Glu  
180 185 190

Leu Asp Thr Arg Arg Ser Gly Arg Gln Gln Leu Gln Ser Ile Lys Gly  
 195 200 205  
 Thr Lys Leu Thr Ile Thr Gln Ala Val Thr Thr Thr Thr Thr Trp Arg  
 210 215 220  
 Pro Ser Ser Thr Thr Thr Ile Ala Gly Leu Arg Val Thr Glu Ser Lys  
 225 230 235 240  
 Gly His Ser Glu Ser Trp His Leu Ser Leu Asp Thr Ala Ile Arg Val  
 245 250 255  
 Ala Leu Ala Val Ala Val Leu Lys Thr Val Ile Leu Gly Leu Leu Cys  
 260 265 270  
 Leu Leu Leu Leu Trp Trp Arg Arg Arg Lys Gly Ser Arg Ala Pro Ser  
 275 280 285  
 Ser Asp Phe  
 290

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 293 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Met Thr Val Ser Gln Arg Phe Gln Leu Ser Asn Ser Gly Pro Asn Ser  
 1 5 10 15  
 Thr Ile Lys Met Lys Ile Ala Leu Arg Val Leu His Leu Glu Lys Arg  
 20 25 30  
 Glu Arg Pro Pro Asp His Gln His Ser Ala Gln Val Lys Arg Pro Ser  
 35 40 45  
 Val Ser Lys Glu Gly Arg Lys Thr Ser Ile Lys Ser His Met Ser Gly  
 50 55 60  
 Ser Pro Gly Pro Gly Gly Ser Asn Thr Ala Pro Ser Thr Pro Val Ile

65                      70                      75                      80  
 Gly Gly Ser Asp Lys Pro Gly Met Glu Glu Lys Ala Gln Pro Pro Glu  
                                  85                      90                      95  
 Ala Gly Pro Gln Gly Leu His Asp Leu Gly Arg Ser Ser Ser Ser Leu  
                                  100                      105                      110  
 Leu Ala Ser Pro Gly His Ile Ser Val Lys Glu Pro Thr Pro Ser Ile  
                                  115                      120                      125  
 Ala Ser Asp Ile Ser Leu Pro Ile Ala Thr Gln Glu Leu Arg Gln Arg  
                                  130                      135                      140  
 Leu Arg Gln Leu Glu Asn Gly Thr Thr Leu Gly Gln Ser Pro Leu Gly  
 145                                   150                      155                      160  
 Gln Ile Gln Leu Thr Ile Arg His Ser Ser Gln Arg Asn Lys Leu Ile  
                                  165                      170                      175  
 Val Val Val His Ala Cys Arg Asn Leu Ile Ala Phe Ser Glu Asp Gly  
                                  180                      185                      190  
 Ser Asp Pro Tyr Val Arg Met Tyr Leu Leu Pro Asp Lys Arg Arg Ser  
                                  195                      200                      205  
 Gly Arg Arg Lys Thr His Val Ser Lys Lys Thr Leu Asn Pro Val Phe  
                                  210                      215                      220  
 Asp Gln Ser Phe Asp Phe Ser Val Ser Leu Pro Glu Val Gln Arg Arg  
 225                                   230                      235                      240  
 Thr Leu Asp Val Ala Val Lys Asn Ser Gly Gly Phe Leu Ser Lys Asp  
                                  245                      250                      255  
 Lys Gly Leu Leu Gly Lys Val Leu Val Ala Leu Ala Ser Glu Glu Leu  
                                  260                      265                      270  
 Ala Lys Gly Trp Thr Gln Trp Tyr Asp Leu Thr Glu Asp Gly Thr Arg  
                                  275                      280                      285  
 Pro Gln Ala Met Thr  
 290

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 206 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Met Glu Arg Arg His Pro Val Cys Ser Gly Thr Cys Gln Pro Thr Gln  
1 5 10 15  
Phe Arg Cys Ser Asn Gly Cys Cys Ile Asp Ser Phe Leu Glu Cys Asp  
20 25 30  
Asp Thr Pro Asn Cys Pro Asp Ala Ser Asp Glu Ala Ala Cys Glu Lys  
35 40 45  
Tyr Thr Ser Gly Phe Asp Glu Leu Gln Arg Ile His Phe Pro Ser Asp  
50 55 60  
Lys Gly His Cys Val Asp Leu Pro Asp Thr Gly Leu Cys Lys Glu Ser  
65 70 75 80  
Ile Pro Arg Trp Tyr Tyr Asn Pro Phe Ser Glu His Cys Ala Arg Phe  
85 90 95  
Thr Tyr Gly Gly Cys Tyr Gly Asn Lys Asn Asn Phe Glu Glu Glu Gln  
100 105 110  
Gln Cys Leu Glu Ser Cys Arg Gly Ile Ser Lys Lys Asp Val Phe Gly  
115 120 125  
Leu Arg Arg Glu Ile Pro Ile Pro Ser Thr Gly Ser Val Glu Met Ala  
130 135 140  
Val Ala Val Phe Leu Val Ile Cys Ile Val Val Val Ala Ile Leu  
145 150 155 160  
Gly Tyr Cys Phe Phe Lys Asn Gln Arg Lys Asp Phe His Gly His His  
165 170 175  
His His Pro Pro Pro Thr Pro Ala Ser Ser Thr Val Ser Thr Thr Glu  
180 185 190  
Asp Thr Glu His Leu Val Tyr Asn His Thr Thr Arg Pro Leu  
195 200 205

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 220 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

```

Met Ala Gly Leu Ser Arg Gly Ser Ala Arg Ala Leu Leu Ala Ala Leu
 1           5           10           15
Leu Ala Ser Thr Leu Leu Ala Leu Leu Val Ser Pro Ala Arg Gly Arg
          20           25           30
Gly Gly Arg Asp His Gly Asp Trp Asp Glu Ala Ser Arg Leu Pro Pro
          35           40           45
Leu Pro Pro Arg Glu Asp Ala Ala Arg Val Ala Arg Phe Val Thr His
          50           55           60
Val Ser Asp Trp Gly Ala Leu Ala Thr Ile Ser Thr Leu Glu Ala Val
65           70           75           80
Arg Gly Arg Pro Phe Ala Asp Val Leu Ser Leu Ser Asp Gly Pro Pro
          85           90           95
Gly Ala Gly Ser Gly Val Pro Tyr Phe Tyr Leu Ser Pro Leu Gln Leu
          100          105          110
Ser Val Ser Asn Leu Gln Glu Asn Pro Tyr Ala Thr Leu Thr Met Thr
          115          120          125
Leu Ala Gln Thr Asn Phe Cys Lys Lys His Gly Phe Asp Pro Gln Ser
          130          135          140
Pro Leu Cys Val His Ile Met Leu Ser Gly Thr Val Thr Lys Val Asn
145          150          155          160
Glu Thr Glu Met Asp Ile Ala Lys His Ser Leu Phe Ile Arg His Pro
          165          170          175
Glu Met Lys Thr Trp Pro Ser Ser His Asn Trp Phe Phe Ala Lys Leu
          180          185          190
Asn Ile Thr Asn Ile Trp Val Leu Asp Tyr Phe Gly Gly Pro Lys Ile
          195          200          205
Val Thr Pro Glu Glu Tyr Tyr Asn Val Thr Val Gln

```

210

215

220

## (2) INFORMATION FOR SEQ ID NO:25:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 197 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: None

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

```

Met Asp His His Cys Pro Trp Leu Asn Asn Cys Val Gly His Tyr Asn
 1             5             10             15
His Arg Tyr Phe Phe Ser Phe Cys Phe Phe Met Thr Leu Gly Cys Val
          20             25             30
Tyr Cys Ser Tyr Gly Ser Trp Asp Leu Phe Arg Glu Ala Tyr Ala Ala
          35             40             45
Ile Glu Lys Met Lys Gln Leu Asp Lys Asn Lys Leu Gln Ala Val Ala
          50             55             60
Asn Gln Thr Tyr His Gln Thr Pro Pro Pro Thr Phe Ser Phe Arg Glu
65             70             75             80
Arg Met Thr His Lys Ser Leu Val Tyr Leu Trp Phe Leu Cys Ser Ser
          85             90             95
Val Ala Leu Ala Leu Gly Ala Leu Thr Val Trp His Ala Val Leu Ile
          100            105            110
Ser Arg Gly Glu Thr Ser Ile Glu Arg His Ile Asn Lys Lys Glu Arg
          115            120            125
Arg Arg Leu Gln Ala Lys Gly Arg Val Phe Arg Asn Pro Tyr Asn Tyr
          130            135            140
Gly Cys Leu Asp Asn Trp Lys Val Phe Leu Gly Val Asp Thr Gly Arg
          145            150            155            160
His Trp Leu Thr Arg Val Leu Leu Pro Ser Thr His Leu Pro His Gly
          165            170            175

```

Asn Gly Met Ser Trp Glu Pro Pro Pro Trp Val Thr Ala His Ser Ala  
180 185 190  
Ser Val Met Ala Val  
195

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 451 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Met	Ala	Pro	Leu	Gly	Met	Leu	Leu	Gly	Leu	Leu	Met	Ala	Ala	Cys	Phe
1				5					10					15	
Thr	Phe	Cys	Leu	Ser	His	Gln	Asn	Leu	Lys	Glu	Phe	Ala	Leu	Thr	Asn
			20					25						30	
Pro	Glu	Lys	Ser	Ser	Thr	Lys	Glu	Thr	Glu	Arg	Lys	Glu	Thr	Lys	Ala
			35					40						45	
Glu	Glu	Glu	Leu	Asp	Ala	Glu	Val	Leu	Glu	Val	Phe	His	Pro	Thr	His
			50					55						60	
Glu	Trp	Gln	Ala	Leu	Gln	Pro	Gly	Gln	Ala	Val	Pro	Ala	Gly	Ser	His
65				70				75						80	
Val	Arg	Leu	Asn	Leu	Gln	Thr	Gly	Glu	Arg	Glu	Ala	Lys	Leu	Gln	Tyr
			85					90						95	
Glu	Asp	Lys	Phe	Arg	Asn	Asn	Leu	Lys	Gly	Lys	Arg	Leu	Asp	Ile	Asn
			100					105						110	
Thr	Asn	Thr	Tyr	Thr	Ser	Gln	Asp	Leu	Lys	Ser	Ala	Leu	Ala	Lys	Phe
			115					120						125	
Lys	Glu	Gly	Ala	Glu	Met	Glu	Ser	Ser	Lys	Glu	Asp	Lys	Ala	Arg	Gln
			130					135						140	
Ala	Glu	Val	Lys	Arg	Leu	Phe	Arg	Pro	Ile	Glu	Glu	Leu	Lys	Lys	Asp

145	150	155	160
Phe Asp Glu Leu Asn Val Val Ile Glu Thr Asp Met Gln Ile Met Val			
165	170	175	
Arg Leu Ile Asn Lys Phe Asn Ser Ser Ser Ser Ser Leu Glu Glu Lys			
180	185	190	
Ile Ala Ala Leu Phe Asp Leu Glu Tyr Tyr Val His Gln Met Asp Asn			
195	200	205	
Ala Gln Asp Leu Leu Ser Phe Gly Gly Leu Gln Val Val Ile Asn Gly			
210	215	220	
Leu Asn Ser Thr Glu Pro Leu Val Lys Glu Tyr Ala Ala Phe Val Leu			
225	230	235	240
Gly Ala Ala Phe Ser Ser Asn Pro Lys Val Gln Val Glu Ala Ile Glu			
245	250	255	
Gly Gly Ala Leu Gln Lys Leu Leu Val Ile Leu Ala Thr Glu Gln Pro			
260	265	270	
Leu Thr Ala Lys Lys Lys Val Leu Phe Ala Leu Cys Ser Leu Leu Arg			
275	280	285	
His Phe Pro Tyr Ala Gln Arg Gln Phe Leu Lys Leu Gly Gly Leu Gln			
290	295	300	
Val Leu Arg Thr Leu Val Gln Glu Lys Gly Thr Glu Val Leu Ala Val			
305	310	315	320
Arg Val Val Thr Leu Leu Tyr Asp Leu Val Thr Glu Lys Met Phe Ala			
325	330	335	
Glu Glu Glu Ala Glu Leu Thr Gln Glu Met Ser Pro Glu Lys Leu Gln			
340	345	350	
Gln Tyr Arg Gln Val His Leu Leu Pro Gly Leu Trp Glu Gln Gly Trp			
355	360	365	
Cys Glu Ile Thr Ala His Leu Leu Ala Leu Pro Glu His Asp Ala Arg			
370	375	380	
Glu Lys Val Leu Gln Thr Leu Gly Val Leu Leu Thr Thr Cys Arg Asp			
385	390	395	400
Arg Tyr Arg Gln Asp Pro Gln Leu Gly Arg Thr Leu Ala Ser Leu Gln			
405	410	415	
Ala Glu Tyr Gln Val Leu Ala Ser Leu Glu Leu Gln Asp Gly Glu Asp			
420	425	430	
Glu Gly Tyr Phe Gln Glu Leu Leu Gly Ser Val Asn Ser Leu Leu Lys			





Thr Leu Arg Met Val Ala Val Leu Val Ser Arg Thr Val Gly Pro Thr  
                           165                          170                          175  
 Gln Arg Leu Leu Leu Cys Gly Thr Leu Ala Ala Leu His Met Leu Phe  
                           180                          185                          190  
 Leu Leu Tyr Leu His Phe Ala Tyr His Lys Val Val Glu Gly Ile Leu  
                           195                          200                          205  
 Asp Thr Leu Glu Gly Pro Asn Ile Pro Pro Ile Gln Arg Val Pro Arg  
                           210                          215                          220  
 Asp Ile Pro Ala Met Leu Pro Ala Ala Arg Leu Pro Thr Thr Val Leu  
                           225                          230                          235                          240  
 Asn Ala Thr Ala Lys Ala Val Ala Val Thr Leu Gln Ser His  
                           245                          250

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 221 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Met Gly Ser Glu Asn Glu Ala Leu Asp Leu Ser Met Lys Ser Val Pro  
   1                          5                          10                          15  
 Trp Leu Lys Ala Gly Glu Val Ser Pro Pro Ile Phe Gln Glu Asp Ala  
                           20                          25                          30  
 Ala Leu Asp Leu Ser Val Ala Ala His Arg Lys Ser Glu Pro Pro Pro  
                           35                          40                          45  
 Glu Thr Leu Tyr Asp Ser Gly Ala Ser Val Asp Ser Ser Gly His Thr  
                           50                          55                          60  
 Val Met Glu Lys Leu Pro Ser Gly Met Glu Ile Ser Phe Ala Pro Ala  
   65                          70                          75                          80  
 Thr Ser His Glu Ala Pro Ala Met Met Asp Ser His Ile Ser Ser Ser

85	90	95
Asp Ala Ala Thr Glu Met Leu Ser Gln Pro Asn His Pro Ser Gly Glu		
100	105	110
Val Lys Ala Glu Asn Asn Ile Glu Met Val Gly Glu Ser Gln Ala Ala		
115	120	125
Lys Val Ile Val Ser Val Glu Asp Ala Val Pro Thr Ile Phe Cys Gly		
130	135	140
Lys Ile Lys Gly Leu Ser Gly Val Ser Thr Lys Asn Phe Ser Phe Lys		
145	150	155
Arg Glu Asp Ser Val Leu Gln Gly Tyr Asp Ile Asn Ser Gln Gly Glu		
165	170	175
Glu Ser Met Gly Asn Ala Glu Pro Leu Arg Lys Pro Ile Lys Asn Arg		
180	185	190
Ser Ile Lys Leu Lys Lys Val Asn Ser Gln Glu Val His Met Leu Pro		
195	200	205
Ile Lys Lys Gln Arg Leu Ala Thr Phe Phe Pro Arg Lys		
210	215	220

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 266 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Met Val Lys Val Thr Phe Asn Ser Ala Leu Ala Gln Lys Glu Ala Lys			
1	5	10	15
Lys Asp Glu Pro Lys Ser Gly Glu Glu Ala Leu Ile Ile Pro Pro Asp			
20	25	30	
Ala Val Ala Val Asp Cys Lys Asp Pro Asp Asp Val Val Pro Val Gly			
35	40	45	

Gln	Arg	Arg	Ala	Trp	Cys	Trp	Cys	Met	Cys	Phe	Gly	Leu	Ala	Phe	Met
50					55						60				
Leu	Ala	Gly	Val	Ile	Leu	Gly	Gly	Ala	Tyr	Leu	Tyr	Lys	Tyr	Phe	Ala
65				70				75						80	
Leu	Gln	Pro	Asp	Asp	Val	Tyr	Tyr	Cys	Gly	Ile	Lys	Tyr	Ile	Lys	Asp
			85					90					95		
Asp	Val	Ile	Leu	Asn	Glu	Pro	Ser	Ala	Asp	Ala	Pro	Ala	Ala	Leu	Tyr
		100						105					110		
Gln	Thr	Ile	Glu	Glu	Asn	Ile	Lys	Ile	Phe	Glu	Glu	Glu	Glu	Val	Glu
		115					120					125			
Phe	Ile	Ser	Val	Pro	Val	Pro	Glu	Phe	Ala	Asp	Ser	Asp	Pro	Ala	Asn
		130				135					140				
Ile	Val	His	Asp	Phe	Asn	Lys	Lys	Leu	Thr	Ala	Tyr	Leu	Asp	Leu	Asn
145				150				155						160	
Leu	Asp	Lys	Cys	Tyr	Val	Ile	Pro	Leu	Asn	Thr	Ser	Ile	Val	Met	Pro
			165					170					175		
Pro	Arg	Asn	Leu	Leu	Glu	Leu	Leu	Ile	Asn	Ile	Lys	Ala	Gly	Thr	Tyr
		180						185					190		
Leu	Pro	Gln	Ser	Tyr	Leu	Ile	His	Glu	His	Met	Val	Ile	Thr	Asp	Arg
		195					200						205		
Ile	Glu	Asn	Ile	Asp	His	Leu	Gly	Phe	Phe	Ile	Tyr	Arg	Leu	Cys	His
		210					215				220				
Asp	Lys	Glu	Thr	Tyr	Lys	Leu	Gln	Arg	Arg	Glu	Thr	Ile	Lys	Gly	Ile
225				230				235						240	
Gln	Lys	Arg	Glu	Ala	Ser	Asn	Cys	Phe	Ala	Ile	Arg	His	Phe	Glu	Asn
			245					250					255		
Lys	Phe	Ala	Val	Glu	Thr	Leu	Ile	Cys	Ser						
		260						265							

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 251 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Met Pro Thr Gly Asp Phe Asp Ser Lys Pro Ser Trp Ala Asp Gln Val  
1 5 10 15  
Glu Glu Glu Gly Glu Asp Asp Lys Cys Val Thr Ser Glu Leu Leu Lys  
20 25 30  
Gly Ile Pro Leu Ala Thr Gly Asp Thr Ser Pro Glu Pro Glu Leu Leu  
35 40 45  
Pro Gly Ala Pro Leu Pro Pro Pro Lys Glu Val Ile Asn Gly Asn Ile  
50 55 60  
Lys Thr Val Thr Glu Tyr Lys Ile Asp Glu Asp Gly Lys Lys Phe Lys  
65 70 75 80  
Ile Val Arg Thr Phe Arg Ile Glu Thr Arg Lys Ala Ser Lys Ala Val  
85 90 95  
Ala Arg Arg Lys Asn Trp Lys Lys Phe Gly Asn Ser Glu Phe Asp Pro  
100 105 110  
Pro Gly Pro Asn Val Ala Thr Thr Thr Val Ser Asp Asp Val Ser Met  
115 120 125  
Thr Phe Ile Thr Ser Lys Glu Asp Leu Asn Cys Gln Glu Glu Glu Asp  
130 135 140  
Pro Met Asn Lys Phe Lys Gly Gln Lys Ile Val Ser Cys Arg Ile Cys  
145 150 155 160  
Lys Gly Asp His Trp Thr Thr Arg Cys Pro Tyr Lys Asp Thr Leu Gly  
165 170 175  
Pro Met Gln Lys Glu Leu Ala Glu Gln Leu Gly Leu Ser Thr Gly Glu  
180 185 190  
Lys Glu Lys Leu Pro Gly Glu Leu Glu Pro Val Gln Ala Thr Gln Asn  
195 200 205  
Lys Thr Gly Lys Tyr Val Pro Pro Ser Leu Arg Asp Gly Ala Ser Arg  
210 215 220  
Arg Gly Glu Ser Met Gln Pro Asn Arg Arg Ala Asp Asp Asn Ala Thr  
225 230 235 240  
Ile Arg Val Thr Asn Leu Arg Arg Gly His Ala  
245 250

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 377 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Met Arg Arg Leu Asn Arg Lys Lys Thr Leu Ser Leu Val Lys Glu Leu  
1 5 10 15  
Asp Ala Phe Pro Lys Val Pro Glu Ser Tyr Val Glu Thr Ser Ala Ser  
20 25 30  
Gly Gly Thr Val Ser Leu Ile Ala Phe Thr Thr Met Ala Leu Leu Thr  
35 40 45  
Ile Met Glu Phe Ser Val Tyr Gln Asp Thr Trp Met Lys Tyr Glu Tyr  
50 55 60  
Glu Val Asp Lys Asp Phe Ser Ser Lys Leu Arg Ile Asn Ile Asp Ile  
65 70 75 80  
Thr Val Ala Met Lys Cys Gln Tyr Val Gly Ala Asp Val Leu Asp Leu  
85 90 95  
Ala Glu Thr Met Val Ala Ser Ala Asp Gly Leu Val Tyr Glu Pro Thr  
100 105 110  
Val Phe Asp Leu Ser Pro Gln Gln Lys Glu Trp Gln Arg Met Leu Gln  
115 120 125  
Leu Ile Gln Ser Arg Leu Gln Glu Glu His Ser Leu Gln Asp Val Ile  
130 135 140  
Phe Lys Ser Ala Phe Lys Ser Thr Ser Thr Ala Leu Pro Pro Arg Glu  
145 150 155 160  
Asp Asp Ser Ser Gln Ser Pro Asn Ala Cys Arg Ile His Gly His Leu  
165 170 175  
Tyr Val Asn Lys Val Ala Gly Asn Phe His Ile Thr Val Gly Lys Ala  
180 185 190

Ile	Pro	His	Pro	Arg	Gly	His	Ala	His	Leu	Ala	Ala	Leu	Val	Asn	His				
																195	200	205	
Glu	Ser	Tyr	Asn	Phe	Ser	His	Arg	Ile	Asp	His	Leu	Ser	Phe	Gly	Glu				
																210	215	220	
Leu	Val	Pro	Ala	Ile	Ile	Asn	Pro	Leu	Asp	Gly	Thr	Glu	Lys	Ile	Ala				
																225	230	235	240
Ile	Asp	His	Asn	Gln	Met	Phe	Gln	Tyr	Phe	Ile	Thr	Val	Val	Pro	Thr				
																245	250	255	
Lys	Leu	His	Thr	Tyr	Lys	Ile	Ser	Ala	Asp	Thr	His	Gln	Phe	Ser	Val				
																260	265	270	
Thr	Glu	Arg	Glu	Arg	Ile	Ile	Asn	His	Ala	Ala	Gly	Ser	His	Gly	Val				
																275	280	285	
Ser	Gly	Ile	Phe	Met	Lys	Tyr	Asp	Leu	Ser	Ser	Leu	Met	Val	Thr	Val				
																290	295	300	
Thr	Glu	Glu	His	Met	Pro	Phe	Trp	Gln	Phe	Phe	Val	Arg	Leu	Cys	Gly				
																305	310	315	320
Ile	Val	Gly	Gly	Ile	Phe	Ser	Thr	Thr	Gly	Met	Leu	His	Gly	Ile	Gly				
																325	330	335	
Lys	Phe	Ile	Val	Glu	Ile	Ile	Cys	Cys	Arg	Phe	Arg	Leu	Gly	Ser	Tyr				
																340	345	350	
Lys	Pro	Val	Asn	Ser	Val	Pro	Phe	Glu	Asp	Gly	His	Thr	Asp	Asn	His				
																355	360	365	
Leu	Pro	Leu	Leu	Glu	Asn	Asn	Thr	His											
																370	375		

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 250 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Met	Gly	Ser	Gln	His	Ser	Ala	Ala	Ala	Arg	Pro	Ser	Ser	Cys	Arg	Arg	1	5	10	15
Lys	Gln	Glu	Asp	Asp	Arg	Asp	Gly	Leu	Leu	Ala	Glu	Arg	Glu	Gln	Glu	20	25	30	
Glu	Ala	Ile	Ala	Gln	Phe	Pro	Tyr	Val	Glu	Phe	Thr	Gly	Arg	Asp	Ser	35	40	45	
Ile	Thr	Cys	Leu	Thr	Cys	Gln	Gly	Thr	Gly	Tyr	Ile	Pro	Thr	Glu	Gln	50	55	60	
Val	Asn	Glu	Leu	Val	Ala	Leu	Ile	Pro	His	Ser	Asp	Gln	Arg	Leu	Arg	65	70	75	80
Pro	Gln	Arg	Thr	Lys	Gln	Tyr	Val	Leu	Leu	Ser	Ile	Leu	Leu	Cys	Leu	85	90	95	
Leu	Ala	Ser	Gly	Leu	Val	Val	Phe	Phe	Leu	Phe	Pro	His	Ser	Val	Leu	100	105	110	
Val	Asp	Asp	Asp	Gly	Ile	Lys	Val	Val	Lys	Val	Thr	Phe	Asn	Lys	Gln	115	120	125	
Asp	Ser	Leu	Val	Ile	Leu	Thr	Ile	Met	Ala	Thr	Leu	Lys	Ile	Arg	Asn	130	135	140	
Ser	Asn	Phe	Tyr	Thr	Val	Ala	Val	Thr	Ser	Leu	Ser	Ser	Gln	Ile	Gln	145	150	155	160
Tyr	Met	Asn	Thr	Val	Val	Ser	Thr	Tyr	Val	Thr	Thr	Asn	Val	Ser	Leu	165	170	175	
Ile	Pro	Pro	Arg	Ser	Glu	Gln	Leu	Val	Asn	Phe	Thr	Gly	Lys	Ala	Glu	180	185	190	
Met	Gly	Gly	Pro	Phe	Ser	Tyr	Val	Tyr	Phe	Phe	Cys	Thr	Val	Pro	Glu	195	200	205	
Ile	Leu	Val	His	Asn	Ile	Val	Ile	Phe	Met	Arg	Thr	Ser	Val	Lys	Ile	210	215	220	
Ser	Tyr	Ile	Gly	Leu	Met	Thr	Gln	Ser	Ser	Leu	Glu	Thr	His	His	Tyr	225	230	235	240
Val	Asp	Cys	Gly	Gly	Asn	Ser	Thr	Ala	Ile	245	250								

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:



(A) LENGTH: 374 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Met Val Thr Cys Phe His Val Pro Tyr Ser Ala Leu Thr Met Phe Ile  
1 5 10 15  
Ser Thr Glu Gln Thr Glu Arg Asp Ser Ala Thr Ala Tyr Arg Met Thr  
20 25 30  
Val Glu Val Leu Gly Thr Val Leu Gly Thr Ala Ile Gln Gly Gln Ile  
35 40 45  
Val Gly Gln Ala Asp Thr Pro Cys Phe Gln Asp Leu Asn Ser Ser Thr  
50 55 60  
Val Ala Ser Gln Ser Ala Asn His Thr His Gly Thr Thr Ser His Arg  
65 70 75 80  
Glu Thr Gln Lys Ala Tyr Leu Leu Ala Ala Gly Val Ile Val Cys Ile  
85 90 95  
Tyr Ile Ile Cys Ala Val Ile Leu Ile Leu Gly Val Arg Glu Gln Arg  
100 105 110  
Glu Pro Tyr Glu Ala Gln Gln Ser Glu Pro Ile Ala Tyr Phe Arg Gly  
115 120 125  
Leu Arg Leu Val Met Ser His Gly Pro Tyr Ile Lys Leu Ile Thr Gly  
130 135 140  
Phe Leu Phe Thr Ser Leu Ala Phe Met Leu Val Glu Gly Asn Phe Val  
145 150 155 160  
Leu Phe Cys Thr Tyr Thr Leu Gly Phe Arg Asn Glu Phe Gln Asn Leu  
165 170 175  
Leu Leu Ala Ile Met Leu Ser Ala Thr Leu Thr Ile Pro Ile Trp Gln  
180 185 190  
Trp Phe Leu Thr Arg Phe Gly Lys Lys Thr Ala Val Tyr Val Gly Ile  
195 200 205  
Ser Ser Ala Val Pro Phe Leu Ile Leu Val Ala Leu Met Glu Ser Asn



Gly Gln Val Leu Ala Gly Arg Ala Arg Arg Leu Leu Leu Gln Phe Gly  
 20 25 30  
 Val Leu Phe Cys Thr Ile Leu Leu Leu Leu Trp Val Ser Val Phe Leu  
 35 40 45  
 Tyr Gly Ser Phe Tyr Tyr Ser Tyr Met Pro Thr Val Ser His Leu Ser  
 50 55 60  
 Pro Val His Phe Tyr Tyr Arg Thr Asp Cys Asp Ser Ser Thr Thr Ser  
 65 70 75 80  
 Leu Cys Ser Phe Pro Val Ala Asn Val Ser Leu Thr Lys Gly Gly Arg  
 85 90 95  
 Asp Arg Val Leu Met Tyr Gly Gln Pro Tyr Arg Val Thr Leu Glu Leu  
 100 105 110  
 Glu Leu Pro Glu Ser Pro Val Asn Gln Asp Leu Gly Met Phe Leu Val  
 115 120 125  
 Thr Ile Ser Cys Tyr Thr Arg Gly Gly Arg Ile Ile Ser Thr Ser Ser  
 130 135 140  
 Arg Ser Val Met Leu His Tyr Arg Ser Asp Leu Leu Gln Met Leu Asp  
 145 150 155 160  
 Thr Leu Val Phe Ser Ser Leu Leu Leu Phe Gly Phe Ala Glu Gln Lys  
 165 170 175  
 Gln Leu Leu Glu Val Glu Leu Tyr Ala Asp Tyr Arg Glu Asn Ser Tyr  
 180 185 190  
 Val Pro Thr Thr Gly Ala Ile Ile Glu Ile His Ser Lys Arg Ile Gln  
 195 200 205  
 Leu Tyr Gly Ala Tyr Leu Arg Ile His Ala His Phe Thr Gly Leu Arg  
 210 215 220  
 Tyr Leu Leu Tyr Asn Phe Pro Met Thr Cys Ala Phe Ile Gly Val Ala  
 225 230 235 240  
 Ser Asn Phe Thr Phe Leu Ser Val Ile Val Leu Phe Ser Tyr Met Gln  
 245 250 255  
 Trp Val Trp Gly Gly Ile Trp Pro Arg His Arg Phe Ser Leu Gln Val  
 260 265 270  
 Asn Ile Arg Lys Arg Asp Asn Ser Arg Lys Glu Val Gln Arg Arg Ile  
 275 280 285  
 Ser Ala His Gln Pro Gly Pro Glu Gly Gln Glu Glu Ser Thr Pro Gln  
 290 295 300

[illegible]

(i) SEQUENCE CHARACTERISTICS:

- (ii) MOLECULE TYPE: None

## 68

145	150	155	160
Cys Lys Leu Asn Ile Leu Lys Ala Leu Gly Ile Val Pro Asn Leu Pro			
165	170	175	
Cys Thr Asp Asn Val Ala Phe Asp Met Glu Arg Leu Thr Arg Thr Gln			
180	185	190	
Ala Val Asn Arg Arg Ser Ala Leu Gly Asp Leu Ala Gly Asp Asn Ser			
195	200	205	
Leu Gly Leu Glu Pro Leu Arg Thr Ser Gly Ile Ser Pro Leu Pro Gln			
210	215	220	
Asp Gly Glu Leu Thr Pro Arg Thr Gly Glu Ile Asn Ile Ala Val Thr			
225	230	235	240
Lys Glu Trp Phe Ile Ile Ala Ser Phe Gly Leu Leu Ser Ala Leu Thr			
245	250	255	
Leu Cys Tyr Met Ile Ile Arg Ala Thr Ala Ser Leu Asn Ala Asn Glu			
260	265	270	
Val Glu Trp Phe			
275			

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 210 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

Met Ala Asn Ser Gly Leu Gln Leu Leu Gly Phe Ser Met Ala Leu Leu			
1	5	10	15
Gly Trp Val Gly Leu Val Ala Cys Thr Ala Ile Pro Gln Trp Gln Met			
20	25	30	
Ser Ser Tyr Ala Gly Asp Asn Ile Ile Thr Ala Gln Ala Met Tyr Lys			
35	40	45	

Gly Leu Trp Met Asp Cys Val Thr Gln Ser Thr Gly Met Met Ser Cys  
 50 55 60  
 Lys Met Tyr Asp Ser Val Leu Ala Leu Ser Ala Ala Leu Gln Ala Thr  
 65 70 75 80  
 Arg Ala Leu Met Val Val Ser Leu Val Leu Gly Phe Leu Ala Met Phe  
 85 90 95  
 Val Ala Thr Met Gly Met Lys Cys Thr Arg Cys Gly Gly Asp Asp Lys  
 100 105 110  
 Val Lys Lys Ala Arg Ile Ala Met Gly Gly Gly Ile Ile Phe Ile Val  
 115 120 125  
 Ala Gly Leu Ala Ala Leu Val Ala Cys Ser Trp Tyr Gly His Gln Ile  
 130 135 140  
 Val Thr Asp Phe Tyr Asn Pro Leu Ile Pro Thr Asn Ile Lys Tyr Glu  
 145 150 155 160  
 Phe Gly Pro Ala Ile Phe Ile Gly Trp Ala Gly Ser Ala Leu Val Ile  
 165 170 175  
 Leu Gly Gly Ala Leu Leu Ser Cys Ser Cys Pro Gly Asn Glu Ser Lys  
 180 185 190  
 Ala Gly Tyr Arg Ala Pro Arg Ser Tyr Pro Lys Ser Asn Ser Ser Lys  
 195 200 205  
 Glu Tyr  
 210

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 476 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Met Ile Arg Pro Gln Leu Arg Thr Ala Gly Leu Gly Arg Cys Leu Leu

1	5	10	15
Pro Gly Leu Leu Leu Leu Val Pro Val Leu Trp Ala Gly Ala Glu			
20	25	30	
Lys Leu His Thr Gln Pro Ser Cys Pro Ala Val Cys Gln Pro Thr Arg			
35	40	45	
Cys Pro Ala Leu Pro Thr Cys Ala Leu Gly Thr Thr Pro Val Phe Asp			
50	55	60	
Leu Cys Arg Cys Cys Arg Val Cys Pro Ala Ala Glu Arg Glu Val Cys			
65	70	75	80
Gly Gly Ala Gln Gly Gln Pro Cys Ala Pro Gly Leu Gln Cys Leu Gln			
85	90	95	
Pro Leu Arg Pro Gly Phe Pro Ser Thr Cys Gly Cys Pro Thr Leu Gly			
100	105	110	
Gly Ala Val Cys Gly Ser Asp Arg Arg Thr Tyr Pro Ser Met Cys Ala			
115	120	125	
Leu Arg Ala Glu Asn Arg Ala Ala Arg Arg Leu Gly Lys Val Pro Ala			
130	135	140	
Val Pro Val Gln Trp Gly Asn Cys Gly Asp Thr Gly Thr Arg Ser Ala			
145	150	155	160
Gly Pro Leu Arg Arg Asn Tyr Asn Phe Ile Ala Ala Val Val Glu Lys			
165	170	175	
Val Ala Pro Ser Val Val His Val Gln Leu Trp Gly Arg Leu Leu His			
180	185	190	
Gly Ser Arg Leu Val Pro Val Tyr Ser Gly Ser Gly Phe Ile Val Ser			
195	200	205	
Glu Asp Gly Leu Ile Ile Thr Asn Ala His Val Val Arg Asn Gln Gln			
210	215	220	
Trp Ile Glu Val Val Leu Gln Asn Gly Ala Arg Tyr Glu Ala Val Val			
225	230	235	240
Lys Asp Ile Asp Leu Lys Leu Asp Leu Ala Val Ile Lys Ile Glu Ser			
245	250	255	
Asn Ala Glu Leu Pro Val Leu Met Leu Gly Arg Ser Ser Asp Leu Arg			
260	265	270	
Ala Gly Glu Phe Val Val Ala Leu Gly Ser Pro Phe Ser Leu Gln Asn			
275	280	285	
Thr Ala Thr Ala Gly Ile Val Ser Thr Lys Gln Arg Gly Gly Lys Glu			

290	295	300	
Leu Gly Met Lys Asp Ser Asp Met Asp Tyr Val Gln Ile Asp Ala Thr			
305	310	315	320
Ile Asn Tyr Gly Asn Ser Gly Gly Pro Leu Val Asn Leu Asp Gly Asp			
325	330	335	
Val Ile Gly Val Asn Ser Leu Arg Val Thr Asp Gly Ile Ser Phe Ala			
340	345	350	
Ile Pro Ser Asp Arg Val Arg Gln Phe Leu Ala Glu Tyr His Glu His			
355	360	365	
Gln Met Lys Gly Lys Ala Phe Ser Asn Lys Lys Tyr Leu Gly Leu Gln			
370	375	380	
Met Leu Ser Leu Thr Val Pro Leu Ser Glu Glu Leu Lys Met His Tyr			
385	390	395	400
Pro Asp Phe Pro Asp Val Ser Ser Gly Val Tyr Val Cys Lys Val Val			
405	410	415	
Glu Gly Thr Ala Ala Gln Ser Ser Gly Leu Arg Asp His Asp Val Ile			
420	425	430	
Val Asn Ile Asn Gly Lys Pro Ile Thr Thr Thr Thr Asp Val Val Lys			
435	440	445	
Ala Leu Asp Ser Asp Ser Leu Ser Met Ala Val Leu Arg Gly Lys Asp			
450	455	460	
Asn Leu Leu Leu Thr Val Ile Pro Glu Thr Ile Asn			
465	470	475	

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 266 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: None

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:



Met Val Lys Val Thr Phe Asn Ser Ala Leu Ala Gln Lys Glu Ala Lys  
 1 5 10 15  
 Lys Asp Glu Pro Glu Ser Gly Glu Glu Ala Leu Ile Ile Pro Pro Asp  
 20 25 30  
 Ala Val Ala Val Asp Cys Lys Asp Pro Asp Asp Val Val Pro Val Gly  
 35 40 45  
 Gln Arg Arg Ala Trp Cys Trp Cys Met Cys Phe Gly Leu Ala Phe Met  
 50 55 60  
 Leu Ala Gly Val Ile Leu Gly Gly Ala Tyr Leu Tyr Lys Tyr Phe Ala  
 65 70 75 80  
 Leu Gln Pro Asp Asp Val Tyr Tyr Cys Gly Ile Lys Tyr Ile Lys Asp  
 85 90 95  
 Asp Val Ile Leu Asn Glu Pro Ser Ala Asp Ala Pro Ala Ala Leu Tyr  
 100 105 110  
 Gln Thr Ile Glu Glu Asn Ile Lys Ile Phe Glu Glu Glu Glu Val Glu  
 115 120 125  
 Phe Ile Ser Val Pro Val Pro Glu Phe Ala Asp Ser Asp Pro Ala Asn  
 130 135 140  
 Ile Val His Asp Phe Asn Lys Lys Leu Thr Ala Tyr Leu Asp Leu Asn  
 145 150 155 160  
 Leu Asp Lys Cys Tyr Val Ile Pro Leu Asn Thr Ser Ile Val Met Pro  
 165 170 175  
 Pro Arg Asn Leu Leu Glu Leu Leu Ile Asn Ile Lys Ala Gly Thr Tyr  
 180 185 190  
 Leu Pro Gln Ser Tyr Leu Ile His Glu His Met Val Ile Thr Asp Arg  
 195 200 205  
 Ile Glu Asn Ile Asp His Leu Gly Phe Phe Ile Tyr Arg Leu Cys His  
 210 215 220  
 Asp Lys Glu Thr Tyr Lys Leu Gln Arg Arg Glu Thr Ile Lys Gly Ile  
 225 230 235 240  
 Gln Lys Arg Glu Ala Ser Asn Cys Phe Ala Ile Arg His Phe Glu Asn  
 245 250 255  
 Lys Phe Ala Val Glu Thr Leu Ile Cys Ser  
 260 265